



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 393 745 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
03.03.2004 Bulletin 2004/10

(51) Int Cl.7: **A61K 39/39**, C07H 21/00,
A61K 31/70

(21) Application number: **03017211.8**

(22) Date of filing: **29.07.2003**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**
Designated Extension States:
AL LT LV MK

- **Kandimalla, Ekambar**
Southboro, MA 01772-1981 (US)
- **Yu, Dong**
Westboro, MA 01581-1145 (US)
- **Bhagat, Lakshmi**
Brookline, MA 02446 (US)

(30) Priority: **29.07.2002 US 399181 P**
29.07.2002 US 399287 P
29.07.2002 US 399302 P
29.07.2002 US 399343 P
29.07.2002 US 399344 P

(74) Representative: **Grund, Martin, Dr. et al**
Dr. Volker Vossius
Patent- & Rechtsanwaltskanzlei
Gelbelstrasse 6
81679 München (DE)

(71) Applicant: **HYBRIDON, INC.**
Cambridge, Massachusetts 02139 (US)

Remarks:
Amended claims in accordance with Rule 86 (2)
EPC.

(72) Inventors:
• **Agrawal, Sudhir**
Shrewsbury, MA 01545-5456 (US)

(54) **Modulation of immunostimulatory properties of oligonucleotide-based compounds by optimal presentation of 5'ends**

(57) The invention relates to the therapeutic use of oligonucleotides as immunostimulatory agents in immunotherapy applications. More particularly, the invention provides immunomers for use in methods for generating an immune response or for treating a patient in need of immunostimulation. The immunomers of the invention

comprise at least two oligonucleotides linked at their 3' ends, internucleoside linkages or functionalized nucleobase or sugar to a non-nucleotidic linker, at least one of the oligonucleotides being an immunostimulatory oligonucleotide and having an accessible 5' end.

BEST AVAILABLE COPY

EP 1 393 745 A1

Description

BACKGROUND OF THE INVENTION5 **Field of the Invention**

[0001] The invention relates to immunology and immunotherapy applications using oligonucleotides as immunostimulatory agents.

10 **Summary of the Related Art**

[0002] Oligonucleotides have become indispensable tools in modern molecular biology, being used in a wide variety of techniques, ranging from diagnostic probing methods to PCR to antisense inhibition of gene expression and immunotherapy applications. This widespread use of oligonucleotides has led to an increasing demand for rapid, inexpensive and efficient methods for synthesizing oligonucleotides.

[0003] The synthesis of oligonucleotides for antisense and diagnostic applications can now be routinely accomplished. See, e.g., *Methods in Molecular Biology*, Vol. 20: *Protocols for Oligonucleotides and Analogs* pp. 165-189 (S. Agrawal, ed., Humana Press, 1993); *Oligonucleotides and Analogues, A Practical Approach*, pp. 87-108 (F. Eckstein, ed., 1991); and Uhlmann and Peyman, *supra*; Agrawal and Iyer, *Curr. Op. in Biotech.* 6:12 (1995); and *Antisense Research and Applications* (Cooke and Lebleu, eds., CRC Press, Boca Raton, 1993). Early synthetic approaches included phosphodiester and phosphotriester chemistries. For example, Khorana et al., *J. Molec. Biol.* 72:209 (1972) discloses phosphodiester chemistry for oligonucleotide synthesis. Reese, *Tetrahedron Lett.* 34:3143-3179 (1978), discloses phosphotriester chemistry for synthesis of oligonucleotides and polynucleotides. These early approaches have largely given way to the more efficient phosphoramidite and H-phosphonate approaches to synthesis. For example, Beaucage and Caruthers, *Tetrahedron Lett.* 22:1859-1862 (1981), discloses the use of deoxyribonucleoside phosphoramidites in polynucleotide synthesis. Agrawal and Zamecnik, U.S. Patent No. 5,149,798 (1992), discloses optimized synthesis of oligonucleotides by the H-phosphonate approach. Both of these modern approaches have been used to synthesize oligonucleotides having a variety of modified internucleotide linkages. Agrawal and Goodchild, *Tetrahedron Lett.* 28:3539-3542 (1987), teaches synthesis of oligonucleotide methylphosphonates using phosphoramidite chemistry. Connolly et al., *Biochem.* 23:3443 (1984), discloses synthesis of oligonucleotide phosphorothioates using phosphoramidite chemistry. Jager et al., *Biochem.* 27:7237 (1988), discloses synthesis of oligonucleotide phosphoramidates using phosphoramidite chemistry. Agrawal et al., *Proc. Natl. Acad. Sci. (USA)* 85:7079-7083 (1988), discloses synthesis of oligonucleotide phosphoramidates and phosphorothioates using H-phosphonate chemistry.

[0004] More recently, several researchers have demonstrated the validity of the use of oligonucleotides as immunostimulatory agents in immunotherapy applications. The observation that phosphodiester and phosphorothioate oligonucleotides can induce immune stimulation has created interest in developing this side effect as a therapeutic tool. These efforts have focused on phosphorothioate oligonucleotides containing the dinucleotide natural CpG. Kuramoto et al., *Jpn. J. Cancer Res.* 83:1128-1131 (1992) teaches that phosphodiester oligonucleotides containing a palindrome that includes a CpG dinucleotide can induce interferon- α and gamma synthesis and enhance natural killer activity. Krieg et al., *Nature* 371:546-549 (1995) discloses that phosphorothioate CpG-containing oligonucleotides are immunostimulatory. Liang et al., *J. Clin. Invest.* 98:1119-1129 (1996) discloses that such oligonucleotides activate human B cells. Moldoveanu et al., *Vaccine* 16:1216-124 (1998) teaches that CpG-containing phosphorothioate oligonucleotides enhance immune response against influenza virus. McCluskie and Davis, *J. Immunol.* 161:4463-4466 (1998) teaches that CpG-containing oligonucleotides act as potent adjuvants, enhancing immune response against hepatitis B surface antigen.

[0005] Other modifications of CpG-containing phosphorothioate oligonucleotides can also affect their ability to act as modulators of immune response. See, e.g., Zhao et al., *Biochem. Pharmacol.* (1996) 51:173-182; Zhao et al., *Biochem Pharmacol.* (1996) 52:1537-1544; Zhao et al., *Antisense Nucleic Acid Drug Dev.* (1997) 7:495-502; Zhao et al., *Bioorg. Med. Chem. Lett.* (1999) 9:3453-3458; Zhao et al., *Bioorg. Med. Chem. Lett.* (2000) 10:1051-1054; Yu et al., *Bioorg. Med. Chem. Lett.* (2000) 10:2585-2588; Yu et al., *Bioorg. Med. Chem. Lett.* (2001) 11:2263-2267; and Kandimalla et al., *Bioorg. Med. Chem.* (2001) 9:807-813.

[0006] These reports make clear that there remains a need to be able to enhance the immune response caused by immunostimulatory oligonucleotides.

55 **BRIEF SUMMARY OF THE INVENTION**

[0007] The invention provides methods for enhancing the immune response caused by oligonucleotide compounds. The methods according to the invention enable increasing the immunostimulatory effect of immunostimulatory oligo-


nucleotides for immunotherapy applications. The present inventors have surprisingly discovered that modification of an immunostimulatory oligonucleotide to optimally present its 5' end dramatically enhances its immunostimulatory capability. Such an oligonucleotide is referred to herein as an "immunomer."

[0008] In a first aspect, therefore, the invention provides immunomers comprising at least two oligonucleotides linked at their 3' ends, an internucleotide linkage, or a functionalized nucleobase or sugar via a non-nucleotidic linker, at least one of the oligonucleotides being an immunostimulatory oligonucleotide and having an accessible 5' end.

[0009] In one embodiment, the immunomer comprises an immunostimulatory dinucleotide of formula 5'-Pyr-Pur-3', wherein Pyr is a natural or non-natural pyrimidine nucleoside and Pur is a natural or non-natural purine nucleoside.

[0010] In another embodiment, the immunomer comprises an immunostimulatory dinucleotide selected from the group consisting of CpG, C*pG, CpG*, and C*pG*, wherein C is cytidine or 2'-deoxycytidine, C* is 2'-deoxythymidine, arabinocytidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside, G is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, or other non-natural purine nucleoside, and p is an internucleoside linkage selected from the group consisting of phosphodiester, phosphorothioate, and phosphorodithioate. In certain preferred embodiments, the immunostimulatory dinucleotide is not CpG.

[0011] In yet another embodiment, the immunostimulatory oligonucleotide comprises an immunostimulatory domain of formula (III):



wherein:

Y is cytidine, 2'-deoxythymidine, 2'-deoxycytidine, arabinocytidine, 2'-deoxythymidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside;

Z is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, 2'-deoxyinosine, or other non-natural purine nucleoside

N1, at each occurrence, is preferably a naturally occurring or a synthetic nucleoside or an immunostimulatory moiety selected from the group consisting of abasic nucleosides, arabinonucleosides, 2'-deoxyuridine, α -deoxyribonucleosides, β -L-deoxyribonucleosides, and nucleosides linked by a phosphodiester or modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleoside linkage being selected from, without limitation, a linker having a length of from about 2 angstroms to about 200 angstroms, C2-C18 alkyl linker, poly(ethylene glycol) linker, 2-aminobutyl-1,3-propanediol linker, glyceryl linker, 2'-5' internucleoside linkage, and phosphorothioate, phosphorodithioate, or methylphosphonate internucleoside linkage;

Nn, at each occurrence, is a naturally occurring nucleoside or an immunostimulatory moiety, preferably selected from the group consisting of abasic nucleosides, arabinonucleosides, 2'-deoxyuridine, α -deoxyribonucleosides, 2'-O-substituted ribonucleosides, and nucleosides linked by a modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleoside linkage being selected from the group consisting of amino linker, 2'-5' internucleoside linkage, and methylphosphonate internucleoside linkage;

provided that at least one N1 or Nn is an immunostimulatory moiety;

wherein n is a number from 0-30;

wherein the 3' end, an internucleotide linkage, or a functionalized nucleobase or sugar is linked directly or via a non-nucleotidic linker to another oligonucleotide, which may or may not be immunostimulatory.

[0012] In a second aspect, the invention provides immunomer conjugates, comprising an immunomer, as described above, and an antigen conjugated to the immunomer at a position other than the accessible 5' end.

[0013] In a third aspect, the invention provides pharmaceutical formulation comprising an immunomer or an immunomer conjugate according to the invention and a physiologically acceptable carrier.

[0014] In a fourth aspect, the invention provides methods for generating an immune response in a vertebrate, such methods comprising administering to the vertebrate an immunomer or immunomer conjugate according to the invention. In some embodiments, the vertebrate is a mammal.

[0015] In a fifth aspect, the invention provides methods for therapeutically treating a patient having a disease or disorder, such methods comprising administering to the patient an immunomer or immunomer conjugate according to the invention. In various embodiments, the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, asthma, allergy, or a disease caused by a pathogen.

5

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] Figure 1 is a schematic representation of representative immunomers of the invention.

[0017] Figure 2 depicts several representative immunomers of the invention.

10 [0018] Figure 3 depicts a group of representative small molecule linkers suitable for linear synthesis of immunomers of the invention.

[0019] Figure 4 depicts a group of representative small molecule linkers suitable for parallel synthesis of immunomers of the invention.

15 [0020] Figure 5 is a synthetic scheme for the linear synthesis of immunomers of the invention. DMTr = 4,4'-dimethoxytrityl; CE = cyanoethyl.

[0021] Figure 6 is a synthetic scheme for the parallel synthesis of immunomers of the invention. DMTr = 4,4'-dimethoxytrityl; CE = cyanoethyl.

20 [0022] Figure 7A is a graphic representation of the induction of IL-12 by immunomers 1-3 in BALB/c mouse spleen cell cultures. These data suggest that Immunomer 2, which has accessible 5'-ends, is a stronger inducer of IL-12 than monomeric Oligo 1, and that Immunomer 3, which does not have accessible 5'-ends, has equal or weaker ability to produce immune stimulation compared with oligo 1.

25 [0023] Figure 7B is a graphic representation of the induction of IL-6 (top to bottom, respectively) by Immunomers 1-3 in BALB/c mouse spleen cells cultures. These data suggest that Immunomer 2, which has accessible 5'-ends, is a stronger inducer of IL-6 than monomeric Oligo 1, and that Immunomer 3, which does not have accessible 5'-ends, has equal or weaker ability to induce immune stimulation compared with Oligo 1.

[0024] Figure 7C is a graphic representation of the induction of IL-10 by Immunomers 1-3 (top to bottom, respectively) in BALB/c mouse spleen cell cultures.

[0025] Figure 8A is a graphic representation of the induction of BALB/c mouse spleen cell proliferation in cell cultures by different concentrations of Immunomers 5 and 6, which have inaccessible and accessible 5'-ends, respectively.

30 [0026] Figure 8B is a graphic representation of BALB/c mouse spleen enlargement by Immunomers 4-6, which have an immunogenic chemical modification in the 5'-flanking sequence of the CpG motif. Again, the immunomer, which has accessible 5'-ends (6), has a greater ability to increase spleen enlargement compared with Immunomer 5, which does not have accessible 5'-end and with monomeric Oligo 4.

35 [0027] Figure 9A is a graphic representation of induction of IL-12 by different concentrations of Oligo 4 and Immunomers 7 and 8 in BALB/c mouse spleen cell cultures.

[0028] Figure 9B is a graphic representation of induction of IL-6 by different concentrations of Oligo 4 and Immunomers 7 and 8 in BALB/c mouse spleen cell cultures.

[0029] Figure 9C is a graphic representation of induction of IL-10 by different concentrations of Oligo 4 and Immunomers 7 and 8 in BALB/c mouse spleen cell cultures.

40 [0030] Figure 10A is a graphic representation of the induction of cell proliferation by Immunomers 14, 15, and 16 in BALB/c mouse spleen cell cultures.

[0031] Figure 10B is a graphic representation of the induction of cell proliferation by IL-12 by different concentrations of Immunomers 14 and 16 in BALB/c mouse spleen cell cultures.

45 [0032] Figure 10C is a graphic representation of the induction of cell proliferation by IL-6 by different concentrations of Immunomers 14 and 16 in BALB/c mouse spleen cell cultures.

[0033] Figure 11A is a graphic representation of the induction of cell proliferation by Oligo 4 and 17 and Immunomers 19 and 20 in BALB/c mouse spleen cell cultures.

[0034] Figure 11B is a graphic representation of the induction of cell proliferation IL-12 by different concentrations of Oligo 4 and 17 and Immunomers 19 and 20 in BALB/c mouse spleen cell cultures.

50 [0035] Figure 11C is a graphic representation of the induction of cell proliferation IL-6 by different concentrations of Oligo 4 and 17 and Immunomers 19 and 20 in BALB/c mouse spleen cell cultures.

[0036] Figure 12 is a graphic representation of BALB/c mouse spleen enlargement using oligonucleotides 4 and immunomers 14, 23, and 24.

55 [0037] Figure 13 is a schematic representation of the 3'-terminal nucleoside of an oligonucleotide, showing that a non-nucleotidic linkage can be attached to the nucleoside at the nucleobase, at the 3' position, or at the 2' position.

[0038] Figure 14 shows the chemical substitutions used in Example 13.

[0039] Figure 15 shows cytokine profiles obtained using the modified oligonucleotides of Example 13.

[0040] Figure 16 shows relative cytokine induction for glycerol linkers compared with amino linkers.

[0041] Figure 17 shows relative cytokine induction for various linkers and linker combinations.

[0042] Figures 18 A-E shows relative nuclease resistance for various PS and PO immunomers and oligonucleotides.

[0043] Figure 19 shows relative cytokine induction for PO immunomers compared with PS immunomers in BALB/c mouse spleen cell cultures.

5 [0044] Figure 20 shows relative cytokine induction for PO immunomers compared with PS immunomers in C3H/HeJ mouse spleen cell cultures.

[0045] Figure 21 shows relative cytokine induction for PO immunomers compared with PS immunomers in C3H/HeJ mouse spleen cell cultures at high concentrations of immunomers.

10 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0046] The invention relates to the therapeutic use of oligonucleotides as immunostimulatory agents for immunotherapy applications. The issued patents, patent applications, and references that are cited herein are hereby incorporated by reference to the same extent as if each was specifically and individually indicated to be incorporated by reference. In the event of inconsistencies between any teaching of any reference cited herein and the present specification, the latter shall prevail for purposes of the invention.

15 [0047] The invention provides methods for enhancing the immune response caused by immunostimulatory compounds used for immunotherapy applications such as, but not limited to, treatment of cancer, autoimmune disorders, asthma, respiratory allergies, food allergies, and bacteria, parasitic, and viral infections in adult and pediatric human and veterinary applications. Thus, the invention further provides compounds having optimal levels of immunostimulatory effect for immunotherapy and methods for making and using such compounds. In addition, immunomers of the invention are useful as adjuvants in combination with DNA vaccines, antibodies, allergens, chemotherapeutic agents, and antisense oligonucleotides.

20 [0048] The present inventors have surprisingly discovered that modification of an immunostimulatory oligonucleotide to optimally present its 5' ends dramatically affects its immunostimulatory capabilities. Such an oligonucleotide is referred to herein as an "immunomer."

[0049] In a first aspect, the invention provides immunomers comprising at least two oligonucleotides linked at their 3' ends, or an internucleoside linkage or a functionalized nucleobase or sugar to a non-nucleotidic linker, at least one of the oligonucleotides being an immunostimulatory oligonucleotide and having an accessible 5' end. As used herein, the term "accessible 5' end" means that the 5' end of the oligonucleotide is sufficiently available such that the factors that recognize and bind to immunomers and stimulate the immune system have access to it. In oligonucleotides having an accessible 5' end, the 5' OH position of the terminal sugar is not covalently linked to more than two nucleoside residues. Optionally, the 5' OH can be linked to a phosphate, phosphorothioate, or phosphorodithioate moiety, an aromatic or aliphatic linker, cholesterol, or another entity which does not interfere with accessibility.

30 [0050] For purposes of the invention, the term "immunomer" refers to any compound comprising at least two oligonucleotides linked at their 3' ends or internucleoside linkages, or functionalized nucleobase or sugar directly or via a non-nucleotidic linker, at least one of the oligonucleotides (in the context of the immunomer) being an immunostimulatory oligonucleotide and having an accessible 5' end, wherein the compound induces an immune response when administered to a vertebrate. In some embodiments, the vertebrate is a mammal, including a human.

40 [0051] In some embodiments, the immunomer comprises two or more immunostimulatory oligonucleotides, (in the context of the immunomer) which may be the same or different. Preferably, each such immunostimulatory oligonucleotide has at least one accessible 5' end.

[0052] In certain embodiments, in addition to the immunostimulatory oligonucleotide(s), the immunomer also comprises at least one oligonucleotide that is complementary to a gene. As used herein, the term "complementary to" means that the oligonucleotide hybridizes under physiological conditions to a region of the gene. In some embodiments, the oligonucleotide downregulates expression of a gene. Such downregulatory oligonucleotides preferably are selected from the group consisting of antisense oligonucleotides, ribozyme oligonucleotides, small inhibitory RNAs and decoy oligonucleotides. As used herein, the term "downregulate a gene" means to inhibit the transcription of a gene or translation of a gene product. Thus, the immunomers according to these embodiments of the invention can be used to target one or more specific disease targets, while also stimulating the immune system.

50 [0053] In certain embodiments, the immunomer includes a ribozyme or a decoy oligonucleotide. As used herein, the term "ribozyme" refers to an oligonucleotide that possesses catalytic activity. Preferably, the ribozyme binds to a specific nucleic acid target and cleaves the target. As used herein, the term "decoy oligonucleotide" refers to an oligonucleotide that binds to a transcription factor in a sequence-specific manner and arrests transcription activity. Preferably, the ribozyme or decoy oligonucleotide exhibits secondary structure, including, without limitation, stem-loop or hairpin structures. In certain embodiments, at least one oligonucleotide comprising poly(I)-poly(dC). In certain embodiments, at least one set of Nn includes a string of 3 to 10 dGs and/or Gs or 2'-substituted ribo or arabino Gs.

55 [0054] For purposes of the invention, the term "oligonucleotide" refers to a polynucleoside formed from a plurality of

linked nucleoside units. Such oligonucleotides can be obtained from existing nucleic acid sources, including genomic or cDNA, but are preferably produced by synthetic methods. In preferred embodiments each nucleoside unit includes a heterocyclic base and a pentofuranosyl, trehalose, arabinose, 2'-deoxy-2'-substituted arabinose, 2'-O-substituted arabinose or hexose sugar group. The nucleoside residues can be coupled to each other by any of the numerous known internucleoside linkages. Such internucleoside linkages include, without limitation, phosphodiester, phosphorothioate, phosphorodithioate, alkylphosphonate, alkylphosphonothioate, phosphotriester, phosphoramidate, siloxane, carbonate, carboalkoxy, acetamidate, carbamate, morpholino, borano, thioether, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphorothioate, and sulfone internucleoside linkages. The term "oligonucleotide" also encompasses polynucleosides having one or more stereospecific internucleoside linkage (e.g., (*R_P*)- or (*S_P*)-phosphorothioate, alkylphosphonate, or phosphotriester linkages). As used herein, the terms "oligonucleotide" and "dinucleotide" are expressly intended to include polynucleosides and dinucleosides having any such internucleoside linkage, whether or not the linkage comprises a phosphate group. In certain preferred embodiments, these internucleoside linkages may be phosphodiester, phosphorothioate, or phosphorodithioate linkages, or combinations thereof.

[0055] In some embodiments, the oligonucleotides each have from about 3 to about 35 nucleoside residues, preferably from about 4 to about 30 nucleoside residues, more preferably from about 4 to about 20 nucleoside residues. In some embodiments, the oligonucleotides have from about 5 to about 18, or from about 5 to about 14, nucleoside residues. As used herein, the term "about" implies that the exact number is not critical. Thus, the number of nucleoside residues in the oligonucleotides is not critical, and oligonucleotides having one or two fewer nucleoside residues, or from one to several additional nucleoside residues are contemplated as equivalents of each of the embodiments described above. In some embodiments, one or more of the oligonucleotides have 11 nucleotides.

[0056] The term "oligonucleotide" also encompasses polynucleosides having additional substituents including, without limitation, protein groups, lipophilic groups, intercalating agents, diamines, folic acid, cholesterol and adamantane. The term "oligonucleotide" also encompasses any other nucleobase containing polymer, including, without limitation, peptide nucleic acids (PNA), peptide nucleic acids with phosphate groups (PHONA), locked nucleic acids (LNA), morpholino-backbone oligonucleotides, and oligonucleotides having backbone sections with alkyl linkers or amino linkers.

[0057] The oligonucleotides of the invention can include naturally occurring nucleosides, modified nucleosides, or mixtures thereof. As used herein, the term "modified nucleoside" is a nucleoside that includes a modified heterocyclic base, a modified sugar moiety, or a combination thereof. In some embodiments, the modified nucleoside is a non-natural pyrimidine or purine nucleoside, as herein described. In some embodiments, the modified nucleoside is a 2'-substituted ribonucleoside an arabinonucleoside or a 2'-deoxy-2'-fluoroarabinoside.

[0058] For purposes of the invention, the term "2'-substituted ribonucleoside" includes ribonucleosides in which the hydroxyl group at the 2' position of the pentose moiety is substituted to produce a 2'-O-substituted ribonucleoside. Preferably, such substitution is with a lower alkyl group containing 1-6 saturated or unsaturated carbon atoms, or with an aryl group having 6-10 carbon atoms, wherein such alkyl, or aryl group may be unsubstituted or may be substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carboalkoxy, or amino groups. Examples of such 2'-O-substituted ribonucleosides include, without limitation 2'-O-methylribonucleosides and 2'-O-methoxyethylribonucleosides.

[0059] The term "2'-substituted ribonucleoside" also includes ribonucleosides in which the 2'-hydroxyl group is replaced with a lower alkyl group containing 1-6 saturated or unsaturated carbon atoms, or with an amino or halo group. Examples of such 2'-substituted ribonucleosides include, without limitation, 2'-amino, 2'-fluoro, 2'-allyl, and 2'-propargyl ribonucleosides.

[0060] The term "oligonucleotide" includes hybrid and chimeric oligonucleotides. A "chimeric oligonucleotide" is an oligonucleotide having more than one type of internucleoside linkage. One preferred example of such a chimeric oligonucleotide is a chimeric oligonucleotide comprising a phosphorothioate, phosphodiester or phosphorodithioate region and non-ionic linkages such as alkylphosphonate or alkylphosphonothioate linkages (see e.g., Pederson *et al.* U. S. Patent Nos. 5,635,377 and 5,366,878).


[0061] A "hybrid oligonucleotide" is an oligonucleotide having more than one type of nucleoside. One preferred example of such a hybrid oligonucleotide comprises a ribonucleotide or 2'-substituted ribonucleotide region, and a deoxyribonucleotide region (see, e.g., Metelev and Agrawal, U.S. Patent No. 5,652,355, 6,346,614 and 6,143,881).

[0062] For purposes of the invention, the term "immunostimulatory oligonucleotide" refers to an oligonucleotide as described above that induces an immune response when administered to a vertebrate, such as a fish, fowl, or mammal. As used herein, the term "mammal" includes, without limitation rats, mice, cats, dogs, horses, cattle, cows, pigs, rabbits, non-human primates, and humans. Useful immunostimulatory oligonucleotides can be found described in Agrawal *et al.*, WO 98/49288, published November 5, 1998; WO 01/12804, published February 22, 2001; WO 01/55370, published August 2, 2001; PCT/US01/13682, filed April 30, 2001; and PCT/US01/30137, filed September 26, 2001. Preferably, the immunostimulatory oligonucleotide comprises at least one phosphodiester, phosphorothioate, or phosphorodithioate internucleoside linkage.

[0063] In some embodiments, the immunostimulatory oligonucleotide comprises an immunostimulatory dinucleotide

of formula 5'-Pyr-Pur-3', wherein Pyr is a natural or synthetic pyrimidine nucleoside and Pur is a natural or synthetic purine nucleoside. As used herein, the term "pyrimidine nucleoside" refers to a nucleoside wherein the base component of the nucleoside is a pyrimidine base. Similarly, the term "purine nucleoside" refers to a nucleoside wherein the base component of the nucleoside is a purine base. For purposes of the invention, a "synthetic" pyrimidine or purine nucleoside includes a non-naturally occurring pyrimidine or purine base, a non-naturally occurring sugar moiety, or a combination thereof.

[0064] Preferred pyrimidine nucleosides according to the invention have the structure (I):



(I)

wherein:

D is a hydrogen bond donor;

D' is selected from the group consisting of hydrogen, hydrogen bond donor, hydrogen bond acceptor, hydrophilic group, hydrophobic group, electron withdrawing group and electron donating group;

A is a hydrogen bond acceptor or a hydrophilic group;

A' is selected from the group consisting of hydrogen bond acceptor, hydrophilic group, hydrophobic group, electron withdrawing group and electron donating group;

X is carbon or nitrogen; and

S' is a pentose or hexose sugar ring, or a non-naturally occurring sugar.


[0065] Preferably, the sugar ring is derivatized with a phosphate moiety, modified phosphate moiety, or other linker moiety suitable for linking the pyrimidine nucleoside to another nucleoside or nucleoside analog.

[0066] Preferred hydrogen bond donors include, without limitation, -NH-, -NH₂, -SH and -OH. Preferred hydrogen bond acceptors include, without limitation, C=O, C=S, and the ring nitrogen atoms of an aromatic heterocycle, e.g., N3 of cytosine.

[0067] In some embodiments, the base moiety in (I) is a non-naturally occurring pyrimidine base. Examples of preferred non-naturally occurring pyrimidine bases include, without limitation, 5-hydroxycytosine, 5-hydroxymethylcytosine, N4-alkylcytosine, preferably N4-ethylcytosine, and 4-thiouracil. However, in some embodiments 5-bromocytosine is specifically excluded.

[0068] In some embodiments, the sugar moiety S' in (I) is a non-naturally occurring sugar moiety. For purposes of the present invention, a "naturally occurring sugar moiety" is a sugar moiety that occurs naturally as part of nucleic acid, e.g., ribose and 2'-deoxyribose, and a "non-naturally occurring sugar moiety" is any sugar that does not occur naturally as part of a nucleic acid, but which can be used in the backbone for an oligonucleotide, e.g., hexose. Arabinose and arabinose derivatives are examples of a preferred sugar moieties.

[0069] Preferred purine nucleoside analogs according to the invention have the structure (II):



(II)

wherein:

D is a hydrogen bond donor;

D' is selected from the group consisting of hydrogen, hydrogen bond donor, and hydrophilic group;

A is a hydrogen bond acceptor or a hydrophilic group;

X is carbon or nitrogen;

each L is independently selected from the group consisting of C, O, N and S; and

S' is a pentose or hexose sugar ring, or a non-naturally occurring sugar.


[0070] Preferably, the sugar ring is derivatized with a phosphate moiety, modified phosphate moiety, or other linker moiety suitable for linking the pyrimidine nucleoside to another nucleoside or nucleoside analog.

[0071] Preferred hydrogen bond donors include, without limitation, -NH-, -NH₂, -SH and -OH. Preferred hydrogen bond acceptors include, without limitation, C=O, C=S, -NO₂ and the ring nitrogen atoms of an aromatic heterocycle, e.g., N1 of guanine.

[0072] In some embodiments, the base moiety in (II) is a non-naturally occurring purine base. Examples of preferred non-naturally occurring purine bases include, without limitation, 6-thioguanine and 7-deazaguanine. In some embodiments, the sugar moiety S' in (II) is a naturally occurring sugar moiety, as described above for structure (I).

[0073] In preferred embodiments, the immunostimulatory dinucleotide is selected from the group consisting of CpG, C*_pG, CpG*, and C*_pG*, wherein C is cytidine or 2'-deoxycytidine, C* is 2'-deoxythymidine, arabinocytidine, 2'-deoxythymidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside, G is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, 2'-deoxyinosine, or other non-natural purine nucleoside, and p is an internucleoside linkage selected from the group consisting of phosphodiester, phosphorothioate, and phosphorodithioate. In certain preferred embodiments, the immunostimulatory dinucleotide is not CpG.

[0074] The immunostimulatory oligonucleotides may include immunostimulatory moieties on one or both sides of the immunostimulatory dinucleotide. Thus, in some embodiments, the immunostimulatory oligonucleotide comprises in immunostimulatory domain of structure (III):



wherein:

Y is cytidine, 2'-deoxythymidine, 2'-deoxycytidine, arabinocytidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-deoxythymidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside;

Z is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, 2'-deoxyinosine, or other

non-natural purine nucleoside;

N1, at each occurrence, is preferably a naturally occurring or a synthetic nucleoside or an immunostimulatory moiety selected from the group consisting of abasic nucleosides, arabinonucleosides, 2'-deoxyuridine, α -deoxyribonucleosides, β -L-deoxyribonucleosides, and nucleosides linked by a phosphodiester or modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleotide linkage being selected from, without limitation, a linker having a length of from about 2 angstroms to about 200 angstroms, C2-C18 alkyl linker, poly(ethylene glycol) linker, 2-aminobutyl-1,3-propanediol linker, glyceryl linker, 2'-5' internucleoside linkage, and phosphorothioate, phosphorodithioate, or methylphosphonate internucleoside linkage;

Nn, at each occurrence, is preferably a naturally occurring nucleoside or an immunostimulatory moiety selected from the group consisting of abasic nucleosides, arabinonucleosides, 2'-deoxyuridine, α -deoxyribonucleosides, 2'-O-substituted ribonucleosides, and nucleosides linked by a modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleotide linkage preferably being selected from the group consisting of amino linker, 2'-5' internucleoside linkage, and methylphosphonate internucleoside linkage; provided that at least one N1 or Nn is an immunostimulatory moiety;

wherein n is a number from 0 to 30; and

wherein the 3' end, an internucleoside linker, or a derivatized nucleobase or sugar is linked directly or via a non-nucleotidic linker to another oligonucleotide, which may or may not be immunostimulatory.

[0075] In some preferred embodiments, YZ is arabinocytidine or 2'-deoxy-2'-substituted arabinocytidine and arabinoguanosine or 2'-deoxy-2'-substituted arabinoguanosine. Preferred immunostimulatory moieties include modifications in the phosphate backbones, including, without limitation, methylphosphonates, methylphosphonothioates, phosphotriesters, phosphothiotriesters, phosphorothioates, phosphorodithioates, triester prodrugs, sulfones, sulfonamides, sulfamates, formacetal, N-methylhydroxylamine, carbonate, carbamate, morpholino, boranophosphonate, phosphoramidates, especially primary amino-phosphoramidates, N3 phosphoramidates and N5 phosphoramidates, and stereospecific linkages (e.g., (Rp)- or (Sp)-phosphorothioate, alkylphosphonate, or phosphotriester linkages).

[0076] Preferred immunostimulatory moieties according to the invention further include nucleosides having sugar modifications, including, without limitation, 2'-substituted pentose sugars including, without limitation, 2'-O-methylribose, 2'-O-methoxyethylribose, 2'-O-propargylribose, and 2'-deoxy-2'-fluororibose; 3'-substituted pentose sugars, including, without limitation, 3'-O-methylribose; 1',2'-dideoxyribose; arabinose; substituted arabinose sugars, including, without limitation, 1'-methylarabinose, 3'-hydroxymethylarabinose, 4'-hydroxymethylarabinose, and 2'-substituted arabinose sugars; hexose sugars, including, without limitation, 1,5-anhydrohexitol; and alpha-anomers. In embodiments in which the modified sugar is a 3'-deoxyribonucleoside or a 3'-O-substituted ribonucleoside, the immunostimulatory moiety is attached to the adjacent nucleoside by way of a 2'-5' internucleoside linkage.

[0077] Preferred immunostimulatory moieties according to the invention further include oligonucleotides having other carbohydrate backbone modifications and replacements, including peptide nucleic acids (PNA), peptide nucleic acids with phosphate groups (PHONA), locked nucleic acids (LNA), morpholino backbone oligonucleotides, and oligonucleotides having backbone linker sections having a length of from about 2 angstroms to about 200 angstroms, including without limitation, alkyl linkers or amino linkers. The alkyl linker may be branched or unbranched, substituted or unsubstituted, and chirally pure or a racemic mixture. Most preferably, such alkyl linkers have from about 2 to about 18 carbon atoms. In some preferred embodiments such alkyl linkers have from about 3 to about 9 carbon atoms. Some alkyl linkers include one or more functional groups selected from the group consisting of hydroxy, amino, thiol, thioether, ether, amide, thioamide, ester, urea, and thioether. Some such functionalized alkyl linkers are poly(ethylene glycol) linkers of formula $-O-(CH_2-CH_2-O-)_n$ ($n = 1-9$). Some other functionalized alkyl linkers are peptides or amino acids.

[0078] Preferred immunostimulatory moieties according to the invention further include DNA isoforms, including, without limitation, β -L-deoxyribonucleosides and α -deoxyribonucleosides. Preferred immunostimulatory moieties according to the invention incorporate 3' modifications, and further include nucleosides having unnatural internucleoside linkage positions, including, without limitation, 2'-5', 2'-2', 3'-3' and 5'-5' linkages.

[0079] Preferred immunostimulatory moieties according to the invention further include nucleosides having modified heterocyclic bases, including, without limitation, 5-hydroxycytosine, 5-hydroxymethylcytosine, N4-alkylcytosine, preferably N4-ethylcytosine, 4-thiouracil, 6-thioguanine, 7-deazaguanine, inosine, nitropyrrole, C5-propynylpyrimidine, and diaminopurines, including, without limitation, 2,6-diaminopurine.

[0080] By way of specific illustration and not by way of limitation, for example, in the immunostimulatory domain of structure (III), a methylphosphonate internucleoside linkage at position N1 or Nn is an immunostimulatory moiety, a linker having a length of from about 2 angstroms to about 200 angstroms, C2-C18 alkyl linker at position X1 is an immunostimulatory moiety, and a β -L-deoxyribonucleoside at position X1 is an immunostimulatory moiety. See Table 1 below for representative positions and structures of immunostimulatory moieties. It is to be understood that reference

to a linker as the immunostimulatory moiety at a specified position means that the nucleoside residue at that position is substituted at its 3'-hydroxyl with the indicated linker, thereby creating a modified internucleoside linkage between that nucleoside residue and the adjacent nucleoside on the 3' side. Similarly, reference to a modified internucleoside linkage as the immunostimulatory moiety at a specified position means that the nucleoside residue at that position is linked to the adjacent nucleoside on the 3' side by way of the recited linkage.

Table 1

Position	TYPICAL IMMUNOSTIMULATORY MOIETIES
N1	Naturally-occurring nucleosides, abasic nucleoside, arabinonucleoside, 2'-deoxyuridine, β -L-deoxyribonucleoside C2-C18 alkyl linker, poly(ethylene glycol) linkage, 2-aminobutyl-1,3-propanediol linker (amino linker), 2'-5' internucleoside linkage, methylphosphonate internucleoside linkage
Nn	Naturally-occurring nucleosides, abasic nucleoside, arabinonucleosides, 2'-deoxyuridine, 2'-O-substituted ribonucleoside, 2'-5' internucleoside linkage, methylphosphonate internucleoside linkage, provided that N1 and N2 cannot both be abasic linkages

[0081] Table 2 shows representative positions and structures of immunostimulatory moieties within an immunostimulatory oligonucleotide having an upstream potentiation domain. As used herein, the term "Spacer 9" refers to a poly(ethylene glycol) linker of formula $-O-(CH_2CH_2-O)_n-$, wherein n is 3. The term "Spacer 18" refers to a poly(ethylene glycol) linker of formula $-O-(CH_2CH_2-O)_n-$, wherein n is 6. As used herein, the term "C2-C18 alkyl linker" refers to a linker of formula $O(CH_2)_q-O-$, where q is an integer from 2 to 18. Accordingly, the terms "C3-linker" and "C3-alkyl linker" refer to a linker of formula $-O-(CH_2)_3-O-$. For each of Spacer 9, Spacer 18, and C2-C18 alkyl linker, the linker is connected to the adjacent nucleosides by way of phosphodiester, phosphorothioate, or phosphorodithioate linkages.

Table 2

Position	TYPICAL IMMUNOSTIMULATORY MOIETY
5' N2	Naturally-occurring nucleosides, 2-aminobutyl-1,3-propanediol linker
5' N1	Naturally-occurring nucleosides, β -L-deoxyribonucleoside, C2-C18 alkyl linker, poly(ethylene glycol), abasic linker, 2-aminobutyl-1,3-propanediol linker
3' N1	Naturally-occurring nucleosides, 1',2'-dideoxyribose, 2'-O-methyl-ribonucleoside, C2-C18 alkyl linker, Spacer 9, Spacer 18
3' N2	Naturally-occurring nucleosides, 1',2'-dideoxyribose, 3'-deoxyribonucleoside, β -L-deoxyribonucleoside, 2'-O-propargyl-ribonucleoside, C2-C18 alkyl linker, Spacer 9, Spacer 18, methylphosphonate internucleoside linkage
3' N3	Naturally-occurring nucleosides, 1',2'-dideoxyribose, C2-C18 alkyl linker, Spacer 9, Spacer 18, methylphosphonate internucleoside linkage, 2'-5' internucleoside linkage, d(G) _n , polyI-polydC
3'N 2+ 3' N 3	1',2'-dideoxyribose, β -L-deoxyribonucleoside, C2-C18 alkyl linker, d(G) _n , polyI-polydC
3'N3+ 3' N 4	2'-O-methoxyethyl-ribonucleoside, methylphosphonate internucleoside linkage, d(G) _n , polyI-polydC
3'N5+ 3' N 6	1',2'-dideoxyribose, C2-C18 alkyl linker, d(G) _n , polyI-polydC
5'N1+ 3' N 3	1',2'-dideoxyribose, d(G) _n , polyI-polydC

[0082] Table 3 shows representative positions and structures of immunostimulatory moieties within an immunostimulatory oligonucleotide having a downstream potentiation domain.

Table 3

Position	TYPICAL IMMUNOSTIMULATORY MOIETY
5' N2	methylphosphonate internucleoside linkage
5' N1	methylphosphonate internucleoside linkage
3' N1	1',2'-dideoxyribose, methylphosphonate internucleoside linkage, 2'-O-methyl

Table 3 (continued)

Position	TYPICAL IMMUNOSTIMULATORY MOIETY
3' N2	1',2'-dideoxyribose, β -L-deoxyribonucleoside, C2-C18 alkyl linker, Spacer 9, Spacer 18, 2-aminobutyl-1,3-propanediol linker, methylphosphonate internucleoside linkage, 2'-O-methyl
3'N3	-deoxyribonucleoside, 3'-O-substituted ribonucleoside, 2'-O-propargyl-ribonucleoside
3'N2 + 3' N3	1',2'-dideoxyribose, β -L-deoxyribonucleoside

[0083] The immunomers according to the invention comprise at least two oligonucleotides linked at their 3' ends or internucleoside linkage or a functionalized nucleobase or sugar via a non-nucleotidic linker. For purposes of the invention, a "non-nucleotidic linker" is any moiety that can be linked to the oligonucleotides by way of covalent or non-covalent linkages. Preferably such linker is from about 2 angstroms to about 200 angstroms in length. Several examples of preferred linkers are set forth below. Non-covalent linkages include, but are not limited to, electrostatic interaction, hydrophobic interactions, π -stacking interactions, and hydrogen bonding. The term "non-nucleotidic linker" is not meant to refer to an internucleoside linkage, as described above, e.g., a phosphodiester, phosphorothioate, or phosphorodithioate functional group, that directly connects the 3'-hydroxyl groups of two nucleosides. For purposes of this invention, such a direct 3'-3' linkage is considered to be a "nucleotidic linkage."

[0084] In some embodiments, the non-nucleotidic linker is a metal, including, without limitation, gold particles. In some other embodiments, the non-nucleotidic linker is a soluble or insoluble biodegradable polymer bead.

[0085] In yet other embodiments, the non-nucleotidic linker is an organic moiety having functional groups that permit attachment to the oligonucleotide. Such attachment preferably is by any stable covalent linkage. As a non-limiting example, the linker may be attached to any suitable position on the nucleoside, as illustrated in Figure 13. In some preferred embodiments, the linker is attached to the 3'-hydroxyl. In such embodiments, the linker preferably comprises a hydroxyl functional group, which preferably is attached to the 3'-hydroxyl by means of a phosphodiester, phosphorothioate, phosphorodithioate or non-phosphate-based linkages.

[0086] In some embodiments, the non-nucleotidic linker is a biomolecule, including, without limitation, polypeptides, antibodies, lipids, antigens, allergens, and oligosaccharides. In some other embodiments, the non-nucleotidic linker is a small molecule. For purposes of the invention, a small molecule is an organic moiety having a molecular weight of less than 1,000 Da. In some embodiments, the small molecule has a molecular weight of less than 750 Da.

[0087] In some embodiments, the small molecule is an aliphatic or aromatic hydrocarbon, either of which optionally can include, either in the linear chain connecting the oligonucleotides or appended to it, one or more functional groups selected from the group consisting of hydroxy, amino, thiol, thioether, ether, amide, thioamide, ester, urea, and thiourea. The small molecule can be cyclic or acyclic. Examples of small molecule linkers include, but are not limited to, amino acids, carbohydrates, cyclodextrins, adamantane, cholesterol, haptens and antibiotics. However, for purposes of describing the non-nucleotidic linker, the term "small molecule" is not intended to include a nucleoside.

[0088] In some embodiments, the small molecule linker is glycerol or a glycerol homolog of the formula $\text{HO}-(\text{CH}_2)_o-\text{CH}(\text{OH})-(\text{CH}_2)_p-\text{OH}$, wherein o and p independently are integers from 1 to about 6, from 1 to about 4, or from 1 to about 3. In some other embodiments, the small molecule linker is a derivative of 1,3-diamino-2-hydroxypropane. Some such derivatives have the formula $\text{HO}-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHC}(\text{O})-(\text{CH}_2)_m-\text{OH}$, wherein m is an integer from 0 to about 10, from 0 to about 6, from 2 to about 6, or from 2 to about 4.

[0089] Some non-nucleotidic linkers according to the invention permit attachment of more than two oligonucleotides, as schematically depicted in Figure 1. For example, the small molecule linker glycerol has three hydroxyl groups to which oligonucleotides may be covalently attached. Some immunomers according to the invention, therefore, comprise more than two oligonucleotides linked at their 3' ends to a non-nucleotidic linker. Some such immunomers comprise at least two immunostimulatory oligonucleotides, each having an accessible 5' end.

[0090] The immunomers of the invention may conveniently be synthesized using an automated synthesizer and phosphoramidite approach as schematically depicted in Figures 5 and 6, and further described in the Examples. In some embodiments, the immunomers are synthesized by a linear synthesis approach (see Figure 5). As used herein, the term "linear synthesis" refers to a synthesis that starts at one end of the immunomer and progresses linearly to the other end. Linear synthesis permits incorporation of either identical or un-identical (in terms of length, base composition and/or chemical modifications incorporated) monomeric units into the immunomers.

[0091] An alternative mode of synthesis is "parallel synthesis", in which synthesis proceeds outward from a central linker moiety (see Figure 6). A solid support attached linker can be used for parallel synthesis, as is described in U.S. Patent No. 5,912,332. Alternatively, a universal solid support (such as phosphate attached controlled pore glass support) can be used.

[0092] Parallel synthesis of immunomers has several advantages over linear synthesis: (1) parallel synthesis permits

the incorporation of identical monomeric units; (2) unlike in linear synthesis, both (or all) the monomeric units are synthesized at the same time, thereby the number of synthetic steps and the time required for the synthesis is the same as that of a monomeric unit; and (3) the reduction in synthetic steps improves purity and yield of the final immunomer product.

5 [0093] At the end of the synthesis by either linear synthesis or parallel synthesis protocols, the immunomers may conveniently be deprotected with concentrated ammonia solution or as recommended by the phosphoramidite supplier, if a modified nucleoside is incorporated. The product immunomer is preferably purified by reversed phase HPLC, detritylated, desalted and dialyzed.

10 [0094] Table 4 shows representative immunomers according to the invention.
Additional immunomers are found described in the Examples.

Table 4. Examples of Immunomer Sequences

15

20

25

30

35

40


45

50

55

Oligo or Immunomer No.	Sequences and Modification (5'-3')
1	5'-GAGAACGCTCGACCTT-3'
2	5'-GAGAACGCTCGACCTT-3'-3'-TTCCAGCTCGCAAGAG-5'
3	3'-TTCCAGCTCGCAAGAG-5'-5'-GAGAACGCTCGACCTT-3'
4	5'-CTATCTGACGTTCTCTGT-3'
5	5'-T-3' $\left[\begin{array}{l} \text{HNCO-C}_4\text{H}_8\text{-5'-CTATLTGACGTTCTCTGT-3'} \\ \text{HNCO-C}_4\text{H}_8\text{-5'-CTATLTGACGTTCTCTGT-3'} \end{array} \right.$
6	$\left. \begin{array}{l} \text{5'-CTATLTGACGTTCTCTGT-3'-C}_4\text{H}_8\text{-CONH} \\ \text{5'-CTATLTGACGTTCTCTGT-3'-C}_4\text{H}_8\text{-CONH} \end{array} \right\} \text{3'-C-5'}$
7	$\left. \begin{array}{l} \text{5'-CTATCTGACGTTCTCTGT-3'-C}_4\text{H}_8\text{-CONH} \\ \text{5'-CTATCTGACGTTCTCTGT-3'-C}_4\text{H}_8\text{-CONH} \end{array} \right\} \text{3'-C-5'}$
8	$\left. \begin{array}{l} \text{5'-CTATCTGACGTTCTCTGT-3'} \\ \text{5'-CTATCTGACGTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
9	$\left. \begin{array}{l} \text{5'-CTATCTGAYGTTCTCTGT-3'} \\ \text{5'-CTATCTGAYGTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
10	$\left. \begin{array}{l} \text{5'-CTATCTGACRTTCTCTGT-3'} \\ \text{5'-CTATCTGACRTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
11	$\left. \begin{array}{l} \text{5'-CTALCTGAYGTTCTCTGT-3'} \\ \text{5'-CTALCTGAYGTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
12	$\left. \begin{array}{l} \text{5'-CTALCTGACRTTCTCTGT-3'} \\ \text{5'-CTALCTGACRTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
13	5'-CTGACGTTCTCTGT-3'
14	$\left. \begin{array}{l} \text{5'-CTGACGTTCTCTGT-3'} \\ \text{5'-CTGACGTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
15	$\left. \begin{array}{l} \text{5'-CTGAYGTTCTCTGT-3'} \\ \text{5'-CTGAYGTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$
16	$\left. \begin{array}{l} \text{5'-CTGACRTTCTCTGT-3'} \\ \text{5'-CTGACRTTCTCTGT-3'} \end{array} \right\} \text{3'-C-5'}$

17	5'-XXTGACGTTCTCTGT-3'
18	5'-XXXTGACGTTCTCTGT-3' } 3'-C-5' 5'-XXXTGACGTTCTCTGT-3' }
19	5'-XXXTGAYGTTCTCTGT-3' } 3'-C-5' 5'-XXXTGAYGTTCTCTGT-3' }
20	5'-XXXTGACRTTCTCTGT-3' } 3'-C-5' 5'-XXXTGACRTTCTCTGT-3' }
21	5'-TCTGACGTTCT-3'
22	5'-XXXTCTGACGTTCT-3' } 3'-C-5' 5'-XXXTCTGACGTTCT-3' }
23	5'-XXXTCTGAYGTTCT-3' } 3'-C-5' 5'-XXXTCTGAYGTTCT-3' }
24	5'-XXXTCTGACRTTCT-3' } 3'-C-5' 5'-XXXTCTGACRTTCT-3' }



L = C3-alkyl linker; X = 1',2'-dideoxyriboside; Y = ^{50H} dC; R = 7-deaza-dG

[0095] In a second aspect, the invention provides immunomer conjugates, comprising an immunomer, as described above, and an antigen conjugated to the immunomer at a position other than the accessible 5' end. In some embodiments, the non-nucleotidic linker comprises an antigen, which is conjugated to the oligonucleotide. In some other embodiments, the antigen is conjugated to the oligonucleotide at a position other than its 3' end. In some embodiments, the antigen produces a vaccine effect.

[0096] The antigen is preferably selected from the group consisting of antigens associated with a pathogen, antigens associated with a cancer, antigens associated with an auto-immune disorder, and antigens associated with other diseases such as, but not limited to, veterinary or pediatric diseases. For purposes of the invention, the term "associated with" means that the antigen is present when the pathogen, cancer, auto-immune disorder, food allergy, respiratory allergy, asthma or other disease is present, but either is not present, or is present in reduced amounts, when the pathogen, cancer, auto-immune disorder, food allergy, respiratory allergy, or disease is absent.

[0097] The immunomer is covalently linked to the antigen, or it is otherwise operatively associated with the antigen. As used herein, the term "operatively associated with" refers to any association that maintains the activity of both immunomer and antigen. Nonlimiting examples of such operative associations include being part of the same liposome or other such delivery vehicle or reagent. In embodiments wherein the immunomer is covalently linked to the antigen, such covalent linkage preferably is at any position on the immunomer other than an accessible 5' end of an immunostimulatory oligonucleotide. For example, the antigen may be attached at an internucleoside linkage or may be attached to the non-nucleotidic linker. Alternatively, the antigen may itself be the non-nucleotidic linker.

[0098] In a third aspect, the invention provides pharmaceutical formulations comprising an immunomer or immunomer conjugate according to the invention and a physiologically acceptable carrier. As used herein, the term "physiologically acceptable" refers to a material that does not interfere with the effectiveness of the immunomer and is compatible with a biological system such as a cell, cell culture, tissue, or organism. Preferably, the biological system is a living organism, such as a vertebrate.

[0099] As used herein, the term "carrier" encompasses any excipient, diluent, filler, salt, buffer, stabilizer, solubilizer, lipid, or other material well known in the art for use in pharmaceutical formulations. It will be understood that the characteristics of the carrier, excipient, or diluent will depend on the route of administration for a particular application. The preparation of pharmaceutically acceptable formulations containing these materials is described in, e.g., *Remington's Pharmaceutical Sciences*, 18th Edition, ed. A. Gennaro, Mack Publishing Co., Easton, PA, 1990.

[0100] In a fourth aspect, the invention provides methods for generating an immune response in a vertebrate, such methods comprising administering to the vertebrate an immunomer or immunomer conjugate according to the invention.

In some embodiments, the vertebrate is a mammal. For purposes of this invention, the term "mammal" is expressly intended to include humans. In preferred embodiments, the immunomer or immunomer conjugate is administered to a vertebrate in need of immunostimulation.

[0101] In the methods according to this aspect of the invention, administration of immunomers can be by any suitable route, including, without limitation, parenteral, oral, sublingual, transdermal, topical, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, by gene gun, dermal patch or in eye drop or mouthwash form. Administration of the therapeutic compositions of immunomers can be carried out using known procedures at dosages and for periods of time effective to reduce symptoms or surrogate markers of the disease. When administered systemically, the therapeutic composition is preferably administered at a sufficient dosage to attain a blood level of immunomer from about 0.0001 micromolar to about 10 micromolar. For localized administration, much lower concentrations than this may be effective, and much higher concentrations may be tolerated. Preferably, a total dosage of immunomer ranges from about 0.001 mg per patient per day to about 200 mg per kg body weight per day. It may be desirable to administer simultaneously, or sequentially a therapeutically effective amount of one or more of the therapeutic compositions of the invention to an individual as a single treatment episode.

[0102] In certain preferred embodiments, immunomers according to the invention are administered in combination with vaccines, antibodies, cytotoxic agents, allergens, antibiotics, antisense oligonucleotides, peptides, proteins, gene therapy vectors, DNA vaccines and/or adjuvants to enhance the specificity or magnitude of the immune response. In these embodiments, the immunomers of the invention can variously act as adjuvants and/or produce direct immunostimulatory effects.

[0103] Either the immunomer or the vaccine, or both, may optionally be linked to an immunogenic protein, such as keyhole limpet hemocyanin (KLH), cholera toxin B subunit, or any other immunogenic carrier protein. Any of the plethora of adjuvants may be used including, without limitation, Freund's complete adjuvant, KLH, monophosphoryl lipid A (MPL), alum, and saponins, including QS-21, imiquimod, R848, or combinations thereof.

[0104] For purposes of this aspect of the invention, the term "in combination with" means in the course of treating the same disease in the same patient, and includes administering the immunomer and/or the vaccine and/or the adjuvant in any order, including simultaneous administration, as well as temporally spaced order of up to several days apart. Such combination treatment may also include more than a single administration of the immunomer, and/or independently the vaccine, and/or independently the adjuvant. The administration of the immunomer and/or vaccine and/or adjuvant may be by the same or different routes.

[0105] The methods according to this aspect of the invention are useful for model studies of the immune system. The methods are also useful for the prophylactic or therapeutic treatment of human or animal disease. For example, the methods are useful for pediatric and veterinary vaccine applications.

[0106] In a fifth aspect, the invention provides methods for therapeutically treating a patient having a disease or disorder, such methods comprising administering to the patient an immunomer or immunomer conjugate according to the invention. In various embodiments, the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, allergy, asthma or a disease caused by a pathogen. Pathogens include bacteria, parasites, fungi, viruses, viroids and prions. Administration is carried out as described for the fourth aspect of the invention.

[0107] For purposes of the invention, the term "allergy" includes, without limitation, food allergies and respiratory allergies. The term "airway inflammation" includes, without limitation, asthma. As used herein, the term "autoimmune disorder" refers to disorders in which "self" proteins undergo attack by the immune system. Such term includes autoimmune asthma.

[0108] In any of the methods according to this aspect of the invention, the immunomer or immunomer conjugate can be administered in combination with any other agent useful for treating the disease or condition that does not diminish the immunostimulatory effect of the immunomer. For example, in the treatment of cancer, it is contemplated that the immunomer or immunomer conjugate may be administered in combination with a chemotherapeutic compound.

[0109] The examples below are intended to further illustrate certain preferred embodiments of the invention, and are not intended to limit the scope of the invention.

EXAMPLES

Example 1: Synthesis of Oligonucleotides Containing Immunomodulatory Moieties

[0110] Oligonucleotides were synthesized on a 1 μ mol scale using an automated DNA synthesizer (Expedite 8909; PerSeptive Biosystems, Framingham, MA), following the linear synthesis or parallel synthesis procedures outlined in Figures 5 and 6.

[0111] Deoxyribonucleoside phosphoramidites were obtained from Applied Biosystems (Foster City, CA). 1',2'-dideoxyribose phosphoramidite, propyl-1-phosphoramidite, 2-deoxyuridine phosphoramidite, 1,3-bis-[5-(4,4'-dimethoxytrityl-

yl)pentylamidy]]-2-propanol phosphoramidite and methyl phosphoramidite were obtained from Glen Research (Sterling, VA). β -L-2'-deoxyribonucleoside phosphoramidite, α -2'-deoxyribonucleoside phosphoramidite, mono-DMT-glycerol phosphoramidite and di-DMT-glycerol phosphoramidite were obtained from ChemGenes (Ashland, MA). (4-Aminobutyl)-1,3-propanediol phosphoramidite was obtained from Clontech (Palo Alto, CA). Arabinocytidine phosphoramidite, arabinoguanosine, arabinothymidine and arabinouridine were obtained from Reliable Pharmaceutical (St. Louis, MO). Arabinoguanosine phosphoramidite, arabinothymidine phosphoramidite and arabinouridine phosphoramidite were synthesized at HybriDon, Inc. (Cambridge, MA) (Noronha et al. (2000) *Biochem.*, 39:7050-7062).

[0112] All nucleoside phosphoramidites were characterized by ^{31}P and ^1H NMR spectra. Modified nucleosides were incorporated at specific sites using normal coupling cycles. After synthesis, oligonucleotides were deprotected using concentrated ammonium hydroxide and purified by reverse phase HPLC, followed by dialysis. Purified oligonucleotides as sodium salt form were lyophilized prior to use. Purity was tested by CGE and MALDI-TOF MS.

Example 2: Analysis of Spleen Cell Proliferation

[0113] *In vitro* analysis of splenocyte proliferation was carried out using standard procedures as described previously (see, e.g., Zhao et al., *Biochem Pharma* 51:173-182 (1996)). The results are shown in Figure 8A. These results demonstrate that at the higher concentrations, Immunomer 6, having two accessible 5' ends results in greater splenocyte proliferation than does Immunomer 5, having no accessible 5' end or Oligonucleotide 4, with a single accessible 5' end. Immunomer 6 also causes greater splenocyte proliferation than the LPS positive control.

Example 3: *In vivo* Splenomegaly Assays

[0114] To test the applicability of the *in vitro* results to an *in vivo* model, selected oligonucleotides were administered to mice and the degree of splenomegaly was measured as an indicator of the level of immunostimulatory activity. A single dose of 5 mg/kg was administered to BALB/c mice (female, 4-6 weeks old, Harlan Sprague Dawley Inc, Baltic, CT) intraperitoneally. The mice were sacrificed 72 hours after oligonucleotide administration, and spleens were harvested and weighed. The results are shown in Figure 8B. These results demonstrate that Immunomer 6, having two accessible 5' ends, has a far greater immunostimulatory effect than do Oligonucleotide 4 or Immunomer 5.

Example 4: Cytokine Analysis

[0115] The secretion of IL-12 and IL-6 in vertebrate cells, preferably BALB/c mouse spleen cells or human PBMC, was measured by sandwich ELISA. The required reagents including cytokine antibodies and cytokine standards were purchased from Pharmingen, San Diego, CA. ELISA plates (Costar) were incubated with appropriate antibodies at 5 $\mu\text{g/mL}$ in PBSN buffer (PBS/0.05% sodium azide, pH 9.6) overnight at 4°C and then blocked with PBS/1% BSA at 37 °C for 30 minutes. Cell culture supernatants and cytokine standards were appropriately diluted with PBS/10% FBS, added to the plates in triplicate, and incubated at 25 °C for 2 hours. Plates were overlaid with 1 $\mu\text{g/mL}$ appropriate biotinylated antibody and incubated at 25 °C for 1.5 hours. The plates were then washed extensively with PBS-T Buffer (PBS/0.05% Tween 20) and further incubated at 25 °C for 1.5 hours after adding streptavidin conjugated peroxidase (Sigma, St. Louis, MO). The plates were developed with Sure Blue™ (Kirkegaard and Perry) chromogenic reagent and the reaction was terminated by adding Stop Solution (Kirkegaard and Perry). The color change was measured on a Ceres 900 HDI Spectrophotometer (Bio-Tek Instruments). The results are shown in Table 5A below.

[0116] Human peripheral blood mononuclear cells (PBMCs) were isolated from peripheral blood of healthy volunteers by Ficoll-Paque density gradient centrifugation (Histopaque-1077, Sigma, St. Louis, MO). Briefly, heparinized blood was layered onto the Histopaque-1077 (equal volume) in a conical centrifuge and centrifuged at 400 x g for 30 minutes at room temperature. The buffy coat, containing the mononuclear cells, was removed carefully and washed twice with isotonic phosphate buffered saline (PBS) by centrifugation at 250 x g for 10 minutes. The resulting cell pellet was then resuspended in RPMI 1640 medium containing L-glutamine (MediaTech, Inc., Herndon, VA) and supplemented with 10% heat inactivated FCS and penicillin-streptomycin (100U/ml). Cells were cultured in 24 well plates for different time periods at 1×10^6 cells/ml/well in the presence or absence of oligonucleotides. At the end of the incubation period, supernatants were harvested and stored frozen at -70 °C until assayed for various cytokines including IL-6 (BD Pharmingen, San Diego, CA), IL-10 (BD Pharmingen), IL-12 (BioSource International, Camarillo, CA), IFN- α (BioSource International) and γ (BD Pharmingen) and TNF- α (BD Pharmingen) by sandwich ELISA. The results are shown in Table 5 below.

[0117] In all instances, the levels of IL-12 and IL-6 in the cell culture supernatants were calculated from the standard curve constructed under the same experimental conditions for IL-12 and IL-6, respectively. The levels of IL-10, IFN-gamma and TNF- α in the cell culture supernatants were calculated from the standard curve constructed under the same experimental conditions for IL-10, IFN-gamma and TNF- α , respectively.

Table 5. Immunomer Structure and Immunostimulatory Activity in Human PBMC Cultures

Oligo No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)		IL-6 (pg/mL)	
			D1	D2	D1	D2
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer (PS)	184	332	3077	5369
26	5'-TCTGTCR ₁ TTCT-3' 5'-TCTGTCR ₁ TTCT-3' \searrow X ₁	11mer (PS)	237	352	3724	4892

Oligo No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-10 (pg/mL)		IFN- γ (pg/mL)	
			D1	D2	D1	D2
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer (PS)	37	88	125	84
26	5'-TCTGTCR ₁ TTCT-3' 5'-TCTGTCR ₁ TTCT-3' \searrow X ₁	11mer (PS)	48	139	251	40


Oligo No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	TNF- α (pg/mL)	
			D1	D2
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer (PS)	537	nt
26	5'-TCTGTCR ₁ TTCT-3' 5'-TCTGTCR ₁ TTCT-3' \searrow X ₁	11mer (PS)	681	nt

D1 and D2 are donors 1 and 2.

Table 5A. Immunomer Structure and Immunostimulatory Activity in BALB/c Mouse Spleen Cell Cultures

Oligo No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			3 µg/mL	10 µg/mL
26	5'-TCTGTCR ₁ TTCT-3' 5'-TCTGTCR ₁ TTCT-3' \searrow X ₁	11mer (PS)	870	10670
27	5'-TCTGTCR ₂ TTCT-3' 5'-TCTGTCR ₂ TTCT-3' \searrow X ₁	11mer (PS)	1441	7664
28	5'-TCTGTY ₂ R ₂ TTCT-3' 5'-TCTGTY ₂ R ₂ TTCT-3' \searrow X ₁	11mer (PS)	1208	1021
29	5'-XXTCTGTCR ₁ TTCT-3' 5'-XXTCTGTCR ₁ TTCT-3' \searrow X ₁	11mer (PS)	162	1013
30	5'-CTGTCR ₂ TTCTCTGT-3' 5'-CTGTCR ₂ TTCTCTGT-3' \searrow X ₁	14mer (PO)	264	251
31	5'-CTGTY ₂ R ₂ TTCTCTGT-3' 5'-CTGTY ₂ R ₂ TTCTCTGT-3' \searrow X ₁	14mer (PO)	149	119
32	5'-TCTGACR ₁ TTCT-3' 5'-TCTGACR ₁ TTCT-3' \searrow X ₁	11mer (PS)	2520	9699
33	5'-XXTCTGACR ₁ TTCT-3' 5'-XXTCTGACR ₁ TTCT-3' \searrow X ₁	11mer (PS)	2214	16881
34	5'-TCTGACR ₂ TTCT-3' 5'-TCTGACR ₂ TTCT-3' \searrow X ₁	11mer (PS)	3945	10766
35	5'-TCTGAY ₂ R ₂ TTCT-3' 5'-TCTGAY ₂ R ₂ TTCT-3' \searrow X ₁	11mer (PS)	2573	19411
36	5'-CTGAY ₂ GTTCTCTGT-3' 5'-CTGAY ₂ GTTCTCTGT-3' \searrow X ₁	14mer (PO)	2699	408
37	5'-CTGACR ₂ TTCTCTGT-3' 5'-CTGACR ₂ TTCTCTGT-3' \searrow X ₁	14mer (PO)	839	85
38	5'-CTGAY ₂ R ₂ TTCTCTGT-3' 5'-CTGAY ₂ R ₂ TTCTCTGT-3' \searrow X ₁	14mer (PO)	143	160

Normal phase represents a phosphorothioate linkage; Italic phase represents a phosphodiester linkage.



[0118] In addition, the results shown in Figures 7A-C demonstrate that Oligonucleotide 2, with two accessible 5' ends elevates IL-12 and IL-6, but not IL-10 at lower concentrations than Oligonucleotides 1 or 3, with one or zero accessible 5' ends, respectively.

Example 5: Effect of Chain Length on Immunostimulatory Activity of Immunomers

[0119] In order to study the effect of length of the oligonucleotide chains, immunomers containing 18, 14, 11, and 8 nucleotides in each chain were synthesized and tested for immunostimulatory activity, as measured by their ability to induce secretion of the cytokines IL-12 and IL-6 in BALB/c mouse spleen cell cultures (Tables 6-8). In this, and all subsequent examples, cytokine assays were carried out in BALB/c spleen cell cultures as described in Example 4.

Table 6. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			@ 0.3 µg/mL	@ 0.3 µg/mL
4	5'-CTATCTGACGTTCTCTGT-3'	18mer	1802	176
39	5'-CTATCTGACGTTCTCTGT-3' 5'-CTATCTGACGTTCTCTGT-3' } 3'-T-5'	18mer	1221	148
40	5'-CTGACGTTCTCTGT-3' 5'-CTGACGTTCTCTGT-3' } 3'-T-5'	14mer	2107	548
41	5'-TCTGACGTTCT-3' 5'-TCTGACGTTCT-3' } 3'-T-5'	11mer	3838	1191
42	5'-GACGTTCT-3' 5'-GACGTTCT-3' } 3'-T-5'	8mer	567	52

Table 7. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			1 µg/mL	1 µg/mL
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer	291	65
43	5'-CTATCTGTCGTTCTCTGT-3' 5'-CTATCTGTCGTTCTCTGT-3' } 3'-T-5'	18mer	430	39
44	5'-CTGTCGTTCTCTGT-3' 5'-CTGTCGTTCTCTGT-3' } 3'-T-5'	14mer	813	59
45	5'-CTGTCGTTCTCT-3' 5'-CTGTCGTTCTCT-3' } 3'-T-5'	12mer	1533	123
46	5'-TCTGTCGTTCT-3' 5'-TCTGTCGTTCT-3' } 3'-T-5'	11mer	2933	505
47	5'-GTCGTTCT-3' 5'-GTCGTTCT-3' } 3'-T-5'	8mer	1086	26
48	5'-GTCGTTCT-3' 5'-GTCGTTCT-3' } 3'-T-5'	7mer	585	34
49	5'-GTCGTT-3' 5'-GTCGTT-3' } 3'-T-5'	6mer	764	76
50	5'-TCGTT-3' 5'-TCGTT-3' } 3'-T-5'	5mer	28	29

Table 8. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			1 µg/mL	1 µg/mL
51	5'-CTCACTTTCGTTCTCTGT-3'	18mer	91	73
52	5'-CTCACTTTCGTTCTCTGT-3' 5'-CTCACTTTCGTTCTCTGT-3' } 3'-T-5'	18mer	502	99
53	5'-CTTTCGTTCTCTGT-3' 5'-CTTTCGTTCTCTGT-3' } 3'-T-5'	14mer	683	119
54	5'-CTTTCGTTCTCT-3' 5'-CTTTCGTTCTCT-3' } 3'-T-5'	12mer	633	102
55	5'-TTCGTTCT-3' 5'-TTCGTTCT-3' } 3'-T-5'	8mer	687	243
56	5'-TCGTTCT-3' 5'-TCGTTCT-3' } 3'-T-5'	7mer	592	1252

[0120] The results suggest that the immunostimulatory activity of immunomers increased as the length of the oligonucleotide chains is decreased from 18-mers to 7-mers. Immunomers having oligonucleotide chain lengths as short as 6-mers or 5-mers showed immunostimulatory activity comparable to that of the 18-mer oligonucleotide with a single 5' end. However, immunomers having oligonucleotide chain lengths as short as 6-mers or 5-mers have increased immunostimulatory activity when the linker is in the length of from about 2 angstroms to about 200 angstroms.

Example 6: Immunostimulatory Activity of Immunomers Containing A Non-Natural Pyrimidine or Non-Natural Purine Nucleoside

[0121] As shown in Tables 9-11, immunostimulatory activity was maintained for immunomers of various lengths having a non-natural pyrimidine nucleoside or non-natural purine nucleoside in the immunostimulatory dinucleotide motif.

Table 9. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			@ 3 µg/mL	@ 3 µg/mL
51	5'-CTCACTTTCGTTCTCTGT-3'	18mer	404	348
57	5'-TCTTTYGTTCT-3' 5'-TCTTTYGTTCT-3' } 3'-T-5'	11mer	591	365
58	5'-TCTTTCRTTCT-3' 5'-TCTTTCRTTCT-3' } 3'-T-5'	11mer	303	283
59	5'-TTYGTTCT-3' 5'-TTYGTTCT-3' } 3'-T-5'	8mer	55	66
60	5'-TTCRTTCT-3' 5'-TTCRTTCT-3' } 3'-T-5'	8mer	242	143





Table 10. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			3 µg/mL	3 µg/mL
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer	379	339
61	5'-TCTGTYGTTCT-3' 5'-TCTGTYGTTCT-3' } 3'-T-5'	11mer	1127	470
62	5'-TCTGTCRTTCT-3' 5'-TCTGTCRTTCT-3' } 3'-T-5'	11mer	787	296
63	5'-GTGTTCT-3' 5'-GTGTTCT-3' } 3'-T-5'	8mer	64	126
64	5'-GTCRTTCT-3' 5'-GTCRTTCT-3' } 3'-T-5'	8mer	246	113







Table 11. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			3 µg/mL	3 µg/mL
4	5'-CTATCTGACGTTCTCTGT-3'	18mer	1176	1892
65	5'-CTATCTGAYGTTCTCTGT-3' 5'-CTATCTGAYGTTCTCTGT-3' } 3'-T-5'	18mer	443	192
66	5'-CTATCTGACRTTCTCTGT-3' 5'-CTATCTGACRTTCTCTGT-3' } 3'-T-5'	18mer	627	464
67	5'-CTGAYGTTCTCTGT-3' 5'-CTGAYGTTCTCTGT-3' } 3'-T-5'	14mer	548	152
68	5'-CTGACRTTCTCTGT-3' 5'-CTGACRTTCTCTGT-3' } 3'-T-5'	14mer	1052	1020
69	5'-TCTGAYGTTCT-3' 5'-TCTGAYGTTCT-3' } 3'-T-5'	11mer	2050	2724
70	5'-TCTGACRTTCT-3' 5'-TCTGACRTTCT-3' } 3'-T-5'	11mer	1780	1741
71	5'-GAYGTTCT-3' 5'-GAYGTTCT-3' } 3'-T-5'	8mer	189	55
72	5'-GACRTTCT-3' 5'-GACRTTCT-3' } 3'-T-5'	8mer	397	212




Example 7: Effect of the Linker on Immunostimulatory Activity

[0122] In order to examine the effect of the length of the linker connecting the two oligonucleotides, immunomers that contained the same oligonucleotides, but different linkers were synthesized and tested for immunostimulatory activity. The results shown in Table 12 suggest that linker length plays a role in the immunostimulatory activity of immunomers. The best immunostimulatory effect was achieved with C3-to C6-alkyl linkers or abasic linkers having interspersed phosphate charges.

Table 12. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			0.3 µg/mL	1 µg/mL
4	5'-CTATCTGACGTTCTCTGT-3'	18mer	257	635
73	5'-CTGACGTTCT-3' \searrow_{X_1} 5'-CTGACGTTCT-3'	10mer	697	1454
74	5'-CTGACGTTCT-3' \searrow_{X_2} 5'-CTGACGTTCT-3'	10mer	1162	669
75	5'-CTGACGTTCT-3' \searrow_{X_3} 5'-CTGACGTTCT-3'	10mer	1074	1375
76	5'-CTGACGTTCT-3' \searrow_{X_4} 5'-CTGACGTTCT-3'	10mer	563	705
77	5'-CTGACGTTCT-3' \searrow_{X_5} 5'-CTGACGTTCT-3'	10mer	264	543
78	5'-CTGACGTTCT-3' \searrow_{X_6} 5'-CTGACGTTCT-3'	10mer	1750	2258
79	5'-CTGACGTTCT-3' $\searrow_{(X_3psX_3)}$ 5'-CTGACGTTCT-3'	10mer	2255	2034
80	5'-CTGACGTTCT-3' $\searrow_{(X_3psX_3psX_3)}$ 5'-CTGACGTTCT-3'	10mer	1493	1197
81	5'-CTGACGTTCT-3' $\searrow_{(X_6psX_6)}$ 5'-CTGACGTTCT-3'	10mer	3625	2642
82	5'-CTGACGTTCT-3' $\searrow_{(X_6psX_6psX_6)}$ 5'-CTGACGTTCT-3'	10mer	4248	2988
83	5'-CTGACGTTCT-3' \searrow_{PO_3S} 5'-CTGACGTTCT-3'	10mer	1241	1964



Example 8: Effect of Oligonucleotide Backbone on Immunostimulatory Activity

[0123] In general, immunostimulatory oligonucleotides that contain natural phosphodiester backbones are less im-

munostimulatory than are the same length oligonucleotides with a phosphorothioate backbones. This lower degree of immunostimulatory activity could be due in part to the rapid degradation of phosphodiester oligonucleotides under experimental conditions. Degradation of oligonucleotides is primarily the result of 3'-exonucleases, which digest the oligonucleotides from the 3' end. The immunomers of this example do not contain a free 3' end. Thus, immunomers with phosphodiester backbones should have a longer half life under experimental conditions than the corresponding monomeric oligonucleotides, and should therefore exhibit improved immunostimulatory activity. The results presented in Table 13 demonstrate this effect, with Immunomers 84 and 85 exhibiting immunostimulatory activity as determined by cytokine induction in BALB/c mouse spleen cell cultures.

Table 13. Immunomer Structure and Immunostimulatory Activity

No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			0.3 µg/mL	1 µg/mL
4	5'-CTATCTGACGTTCTCTGT-3'	18mer	225	1462
84	5'-CTGACGTTCTCTGT-3' 5'-CTGACGTTCTCTGT-3' } 3'-T-5' (PO)	14mer	1551	159
85	5'-LLCTGACGTTCTCTGT-3' 5'-LLCTGACGTTCTCTGT-3' } 3'-T-5' (P)	14mer	466	467

L = C3-Linker

Example 9: Synthesis of Immunomers 73-92

[0124] Oligonucleotides were synthesized on 1 µmol scale using an automated DNA synthesizer (Expedite 8909 PerSeptive Biosystems). Deoxynucleoside phosphoramidites were obtained from Applied Biosystems (Foster City, CA). 7-Deaza-2'-deoxyguanosine phosphoramidite was obtained from Glen Research (Sterling Virginia). 1,3-Bis-DMT-glycerol-CPG was obtained from ChemGenes (Ashland, MA). Modified nucleosides were incorporated into the oligonucleotides at specific site using normal coupling cycles. After the synthesis, oligonucleotides were deprotected using concentrated ammonium hydroxide and purified by reversed-phase HPLC, followed by dialysis. Purified oligonucleotides as sodium salt form were lyophilized prior to use. Purity of oligonucleotides was checked by CGE and MALDI-TOF MS (Bruker Proflex III MALDI-TOF Mass spectrometer).

Example 11

Immunomer Stability

[0125] Oligonucleotides were incubated in PBS containing 10% bovine serum at 37° C for 4, 24 or 48 hours. Intact oligonucleotide was determined by capillary gel electrophoresis. The results are shown in Table 14.

Table 14. Digestion of Oligonucleotides in 10 % Bovine Serum PBS Solution

Oligo No.	Sequences and Modification (5'-3')	CE analysis of oligos (% intact oligo remained after digestion)		
		after 4h	After 24h	after 48h
4	5-CTATCTGACGTTCTCTGT-3'/PS	90.9	71.8	54.7
26	(5'-TCTGTCGTTCT) ₂ S/PS (G=dG ^{deaza})	97.1	91.0	88.1
86	(5'-CTGTCGTTCTCTGT) ₂ S/PO		37.8	22.5
87	(5'-XXCTGTCGTTCTCTGT) ₂ S/PO	73.1	56.8	36.8
88	(5'-UCTGTCGTTCTCTGT) ₂ S/PO		48.4	36.7

X = C3-Linker, U, C = 2'-OMe-ribonucleoside

Example 12: Effect of accessible 5' ends on immunostimulatory activity.

[0126] BALB/c mouse (4-8 weeks) spleen cells were cultured in RPMI complete medium. Murine macrophage-like cells, J774 (American Type Culture Collection, Rockville, MD) were cultured in Dulbecco's modified Eagle's medium supplemented with 10% (v/v) FCS and antibiotics (100 IU/mL of penicillin G/streptomycin). All other culture reagents were purchased from Mediatech (Gaithersburg, MD).

[0127] *ELISAs for IL-12 and IL-6.* BALB/c mouse spleen or J774 cells were plated in 24-well dishes at a density of 5×10^6 or 1×10^6 cells/mL, respectively. The CpG DNA dissolved in TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA) was added to a final concentration of 0.03, 0.1, 0.3, 1.0, 3.0, or 10.0 $\mu\text{g/mL}$ to mouse spleen cell cultures and 1.0, 3.0, or 10.0 $\mu\text{g/mL}$ to J774 cell cultures. The cells were then incubated at 37 °C for 24 hr and the supernatants were collected for ELISA assays. The experiments were performed two or three times for each CpG DNA in triplicate for each concentration.

[0128] The secretion of IL-12 and IL-6 was measured by sandwich ELISA. The required reagents, including cytokine antibodies and standards were purchased from PharMingen. ELISA plates (Costar) were incubated with appropriate antibodies at 5 $\mu\text{g/mL}$ in PBSN buffer (PBS/0.05% sodium azide, pH 9.6) overnight at 4 °C and then blocked with PBS/1% BSA at 37 °C for 30 min. Cell culture supernatants and cytokine standards were appropriately diluted with PBS/1% BSA, added to the plates in triplicate, and incubated at 25 °C for 2 hr. Plates were washed and incubated with 1 $\mu\text{g/mL}$ of appropriate biotinylated antibody and incubated at 25 °C for 1.5 hr. The plates were washed extensively with PBS/0.05% Tween 20 and then further incubated at 25 °C for 1.5 hr after the addition of streptavidine-conjugated peroxidase (Sigma). The plates were developed with Sure Blue™ (Kirkegaard and Perry) chromogenic reagent and the reaction was terminated by adding Stop Solution (Kirkegaard and Perry). The color change was measured on a Ceres 900 HDI Spectrophotometer (Bio-Tek Instruments) at 450 nm. The levels of IL-12 and IL-6 in the cell culture supernatants were calculated from the standard curve constructed under the same experimental conditions for IL-12 and IL-6, respectively.

[0129] The results are shown in Table 15.

Table 15: Phosphorothioate CpG DNA sequences and modifications used in the study^a

CpG DNA #	Sequence	Length	5'-end	3'-end
89	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	19-mer	1	1
90	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-b	19-mer	1	blocked
91	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-3'-g-5'	20-mer	2	blocked
92	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-3'-h-5'	23-mer	2	blocked
93	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-3'-i-5'	27-mer	2	blocked
94	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-3'-j-5'	38-mer	2	blocked
95	b-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	19-mer	blocked	1
96	3'-c-5'-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	20-mer	blocked	2
97	3'-d-5'-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	23-mer	blocked	2
98	3'-e-5'-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	27-mer	blocked	2
99	3'-f-5'-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	38-mer	blocked	2
100	5'-TCCATGAC <u>CG</u> TTCTGATGC-3'-k	19-mer	1	blocked
101	l-5'-TCCATGAC <u>CG</u> TTCTGATGC-3'	19-mer	blocked	1

^a: See Chart I for chemical structures b-l; 5'-CG-3' dinucleotide is shown underlined.

Chart 1




Table 16. Induction of IL-12 and IL-6 secretion by CpG DNA-conjugates in BALB/c mice spleen cell cultures

CpG DNA # ^a	IL-12 (pg/mL)±SD					IL-6 (pg/mL)±SD				
	0.1µg/mL	0.3µg/mL	1.0µg/mL	3.0µg/mL	10.0µg/mL	0.1µg/mL	0.3µg/mL	1.0µg/mL	3.0µg/mL	10.0µg/mL
89	991±121	1820±224	2391±175	3507±127	2615±279	652±48	2858±180	13320±960	18625±1504	17229±1750
90	526±32	2100±175	1499±191	3019±35	3489±162	1387±152	1426±124	5420±370	19096±484	19381±2313
91	1030±11	1348±102	2060±130	3330±130	3582±259	923±22	2542±81	9054±120	14114±179	13693±264
92	1119±159	1726±207	2434±100	2966±204	3215±464	870±146	1905±56	7841±350	17146±1246	15713±693
93	1175±68	2246±124	1812±75	2388±320	2545±202	1152±238	3499±116	7142±467	14064±167	13566±477
94	1087±121	1705±163	1797±141	2522±195	3054±103	1039±105	2043±157	4848±288	15527±224	21021±1427
95	1173±107	2170±155	2132±58	2812±203	3689±94	807±0.5	927±0.5	3344±0.5	10233±0.5	9213±0.5
96	866±51	1564±63	1525±63	2666±97	4030±165	750±63	1643±30	5559±415	11549±251	11060±651
97	227±3	495±96	1007±68	897±15	1355±97	302±18	374±22	1000±68	9106±271	13077±381
98	139±18	211±12	452±22	458±29	1178±237	220±23	235±18	383±35	1706±33	11530±254
99	181±85	282±105	846±165	2082±185	3185±63	467±122	437±85	1697±283	9781±13	11213±294
Medium	86±6					60±12				

^a: See Table 1 for sequences.

Taken together, the current results suggest that an accessible 5'-end of CpG DNA is required for its optimal immunostimulatory activity and smaller groups such as a phosphorothioate, a mononucleotide, or a dinucleotide do not effectively block the accessibility of the 5'-end of CpG DNA to receptors or factors involved in the immunostimulatory pathway. However, the conjugation of molecules as large as fluorescein or larger at the 5'-end of CpG DNA could abrogate immunostimulatory activity. These results have a direct impact on the studies of immunostimulatory activity of CpG DNA-antigen/vaccine/monoclonal antibody (mAb) conjugates. The conjugation of large molecules such as vaccines or mAbs at the 5'-end of a CpG DNA could lead to suboptimal immunostimulatory activity of CpG DNA. The conjugation of functional ligands at the 3'-end of CpG DNA not only contributes to increased nuclease stability but also increased immunostimulatory potency of CpG DNA *in vivo*.

Example 13: Effect of linkers on cytokine secretion

[0130] The following oligonucleotides were synthesized for this study. Each of these modified oligonucleotides can be incorporated into an immunomer.

Table 17. Sequences of CpG DNA showing the position of substitution.

CpG DNA Number	Sequence (5'--->3') ^a
102	CCTACTAGCGTTCTCATC
103	CCTACTAGC2TTCTCATC
104	CCTACT2GCGTTCTCATC
105	CCTA2TAGCGTTCTCATC
106	CCT22TAGCGTTCTCATC
107	22TACTAGCGTTCTCATC
108	CCTACTAGCGT2CTCATC
109	CCTACTAGCGTTC2CATC
110	CCTACTAGCGTTC22ATC
111	CCT6CTAGCGTTCTCATC
112	CCTACTAGCGTTC6CATC
113	CCT7CTAGCGTTCTCATC
114	CCTACTAGCGTTC7CATC
4	CTATCTGACGTTCTCTGT

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

115 CTAT1TGACGTTCTCTGT
 116 CTA1CTGACGTTCTCTGT
 117 CTATCTG2CGTTCTCTGT
 118 CTATC2GACGTTCTCTGT
 119 CTA2CTGACGTTCTCTGT
 120 2222TGACGTTCTCTGT
 121 222TGACGTTCTCTGT
 122 22TGACGTTCTCTGT
 123 2TGACGTTCTCTGT
 124 2TGACGTTCTCTGT
 125 CTAT3TGACGTTCTCTGT
 126 CTA3CTGACGTTCTCTGT
 127 CTA33TGACGTTCTCTGT
 128 33TGACGTTCTCTGT
 129 CTAT4TGACGTTCTCTGT
 130 CTA4CTGACGTTCTCTGT
 131 CTA44TGACGTTCTCTGT
 132 44TGACGTTCTCTGT
 133 CTAT5TGACGTTCTCTGT
 134 CTA5CTGACGTTCTCTGT
 135 CTA55TGACGTTCTCTGT
 136 55TGACGTTCTCTGT
 137 CTA6CTGACGTTCTCTGT
 138 CTATCTGACGTTCTCTGT
 139 CTA7CTGACGTTCTCTGT
 140 CTATCTGACGTTCTCTGT
 141 CTATCTG8CGTTCTCTGT
 142 CTATCT8ACGTTCTCTGT
 143 CTATC8GACGTTCTCTGT
 144 CTAT8TGACGTTCTCTGT
 145 CTA8CTGACGTTCTCTGT
 146 CTATCTGACG8TCTCTGT
 147 CTATCTGACG8TCTCTGT
 148 CTATCTGACGTT8TCTGT
 149 CTATCTGACGTTCT8CTGT
 150 CTATCTG9CGTTCTCTGT
 151 CTATCT9ACGTTCTCTGT
 152 CTA9CTGACGTTCTCTGT
 153 CTATCTGACGT9CTCTGT
 154 CTATCTGACGTTCT9CTGT

^a. See Figure 14 for the chemical structures of substitutions 1-9. All CpG DNAs are phosphorothioate backbone modified.

[0131] To evaluate the optimal linker size for potentiation of immunostimulatory activity, we measured IL-12 and IL-6 secretion induced by modified CpG DNAs in BALB/c mouse spleen cell cultures. All CpG DNAs induced concentration-dependent IL-12 and IL-6 secretion. Figure 15 shows data obtained at 1 µg/mL concentration of selected CpG DNAs, 116, 119, 126, 130, and 134, which had a linker at the fifth nucleotide position in the 5'-flanking sequence to the CpG dinucleotide compared with the parent CpG DNA. The CpG DNAs, which contained C2- (1), C3- (2), and C4-linkers (3), induced secretion of IL-12 production similar to that of the parent CpG DNA 4. The CpG DNA that contained C6 and C9-linkers (4 and 5) at the fifth nucleotide position from CpG dinucleotide in the 5'-flanking sequence induced lower levels of IL-12 secretion than did the parent CpG DNA (Fig. 15), suggesting that substitution of linkers longer than a C4-linker results in the induction of lower levels of IL-12. All five CpG DNAs, which had linkers, induced two to three times higher IL-6 secretion than did the parent CpG DNA. The presence of a linker in these CpG DNAs showed a significant effect on the induction of IL-6 compared with CpG DNAs that did not have a linker. However, we did not observe length-dependent linker effect on IL-6 secretion.

[0132] To examine the effect on immunostimulatory activity of CpG DNA containing ethyleneglycol-linkers, we synthesized CpG DNAs 137 and 138, in which a triethyleneglycol-linker (6) is incorporated at the fifth nucleotide position in the 5'- and at the fourth nucleotide position in the 3'-flanking sequences to the CpG dinucleotide, respectively. Similarly, CpG DNAs 139 and 140 contained a hexaethyleneglycol-linker (7) in the 5'- or the 3'-flanking sequence to the CpG dinucleotide, respectively. All four modified CpG DNAs (137-140) were tested in BALB/c mouse spleen cell cultures for cytokine induction (IL-12, IL-6, and IL-10) in comparison with parent CpG DNA 4. All CpG DNAs induced concentration-dependent cytokine production over the concentration range tested (0.03-10.0 µg/mL) (data not shown). The levels of cytokines induced at 0.3 µg/mL concentration of CpG DNAs 137-140 are shown in Table 18. CpG DNAs 137 and 139, which had an ethyleneglycol-linker in the 5'-flanking sequence induced higher levels of IL-12 (2106±143 and 2066±153 pg/mL) and IL-6 (2362±166 and 2507±66 pg/mL) secretion than did parent CpG DNA 4 (Table 18). At the same concentration, 137 and 139 induced slightly lower levels of IL-10 secretion than did the parent CpG DNA (Table 18). CpG DNA 138, which had a shorter ethyleneglycol-linker (6) in the 3'-flanking sequence induced IL-12 secretion similar to that of the parent CpG DNA, but significantly lower levels of IL-6 and IL-10 (Table 18). CpG DNA 140, which had a longer ethyleneglycol-linker (7) induced significantly lower levels of all three cytokines tested compared with the parent CpG DNA (Table 18).

[0133] Though triethyleneglycol-linker (6) had a chain length similar to that of C9-linker (5), the CpG DNA containing triethyleneglycol-linker had better immunostimulatory activity than did CpG DNA containing C9-linker, as determined by induction of cytokine secretion in spleen cell cultures. These results suggest that the lower immunostimulatory activity observed with CpG DNA containing longer alkyl-linkers (4 and 5) may not be related to their increased length but to their hydrophobic characteristics. This observation prompted us to examine substitution of branched alkyl-linkers containing hydrophilic functional groups on immunostimulatory activity.

Table 18. Cytokine secretion induced by CpG DNAs containing an ethyleneglycol-linker in BALB/c mice spleen cell cultures.

CpG DNA Number	Cytokine, pg/mL		
	IL-12	IL-6	IL-10
4	1887±233	2130±221	86±14
137	2106±143	2362±166	78±21
138	1888±259	1082±25	47±14
139	2066±153	2507±66	73±17
140	1318±162	476±13	25±5
Medium	84±13	33±6	2±1

[0134] To test the effect on immunostimulatory activity of CpG DNA containing branched alkyl-linkers, two branched alkyl-linkers containing a hydroxyl (8) or an amine (9) functional group were incorporated in parent CpG DNA 4 and

the effects on immunostimulatory activity of the resulting modified CpG DNAs (150-154-Table 19) were examined. The data obtained with CpG DNAs 150-154, containing amino-linker 9 at different nucleotide positions, in BALB/c mouse spleen cell cultures (proliferation) and *in vivo* (splenomegaly) are shown in Table 19.

Table 19.

Spleen cell proliferation induced by CpG DNA containing an aminobutyryl propanedoil-linker in BALB/c mice spleen cell cultures and splenomegaly in BALB/c mice.		
CpG DNA Number ^a	Spleen cell proliferation (PI) ^b	Spleen weight (mg) ^c
4	3.7±0.8	121±15
150	2.5±0.6	107±11
151	9.2±0.7	169±16
152	8.8±0.4	220±8
153	7.6±0.7	127±24
154	7.8±0.04	177±12
M/V	1.2±0.3	102±8
LPS	2.8±0.5	ND

[0135] Parent CpG DNA 4 showed a proliferation index of 3.7±0.8 at a concentration of 0.1 µg/mL. At the same concentration, modified CpG DNAs 151-154 containing amino-linker 9 at different positions caused higher spleen cell proliferation than did the parent CpG DNA (Table 19). As observed with other linkers, when the substitution was placed adjacent to CpG dinucleotide (150), a lower proliferation index was noted compared with parent CpG DNA (Table 19), further confirming that the placement of a linker substitution adjacent to CpG dinucleotide has a detrimental effect on immunostimulatory activity. In general, substitution of an amino-linker for 2'-deoxyribonucleoside in the 5'-flanking sequence (151 and 152) resulted in higher spleen cell proliferation than found with the substitution in the 3'-flanking sequence (153 and 154). Similar results were observed in the splenomegaly assay (Table 19), confirming the results observed in spleen cell cultures. Modified CpG DNAs containing glycerol-linker (8) showed immunostimulatory activity similar to or slightly higher than that observed with modified CpG DNA containing amino-linker (9) (data not shown).

[0136] In order to compare the immunostimulatory effects of CpG DNA containing linkers 8 and 9, we selected CpG DNAs 145 and 152, which had substitution in the 5'-flanking sequence and assayed their ability to induce cytokines IL-12 and IL-6 secretion in BALB/c mouse spleen cell cultures. Both CpG DNAs 145 and 152 induced concentration-dependent cytokine secretion. Figure 4 shows the levels of IL-12 and IL-6 induced by 145 and 152 in mouse spleen cell cultures at 0.3 µg/mL concentration compared with parent CpG DNA 4. Both CpG DNAs induced higher levels of IL-12 and IL-6 than did parent CpG DNA 4. CpG DNA containing glycerol-linker (8) induced slightly higher levels of cytokines (especially IL-12) than did CpG DNA containing amino-linker (9) (Figure 16). These results further confirm that the linkers containing hydrophilic groups are more favorable for immunostimulatory activity of CpG DNA.

[0137] We examined two different aspects of multiple linker substitutions in CpG DNA. In one set of experiments, we kept the length of nucleotide sequence to 13-mer and incorporated one to five C3-linker (2) substitutions at the 5'-end (120-124). These modified CpG DNAs permitted us to study the effect of an increase in the length of linkers without causing solubility problems. In the second set of experiments, we incorporated two of the same linker substitutions (3, 4, or 5) in adjacent positions in the 5'-flanking sequence to the CpG dinucleotide to study if there would be any additive effect on immunostimulatory activity.

[0138] Modified CpG DNAs were studied for their ability to induce cytokine production in BALB/c mouse spleen cell cultures in comparison with parent CpG DNA 4. All CpG DNAs induced concentration-dependent cytokine production. The data obtained at 1.0 µg/mL concentration of CpG DNAs is shown in Table 20. In this assay, parent CpG DNA 4 induced 967±28 pg/mL of IL-12, 1593±94 pg/mL of IL-6, and 14±6 pg/mL of IL-10 secretion at 1 µg/mL of concentration. The data presented in Table 20 suggest that as the number of linker substitutions decreased IL-12 induction decreased. However, the induction of lower levels of IL-12 secretion by CpG DNAs 123 and 124 could be the result of the shorter length of CpG DNAs. Our studies with unmodified CpG DNA shorter than 15-nucleotides showed insignificant immunostimulatory activity (data not shown). Neither length nor the number of linker substitutions have a lesser effect on IL-6 secretion. Though IL-10 secretion increased with linker substitutions, the overall IL-10 secretion by these CpG DNAs was minimal.

[0139] CpG DNAs containing two linker substitutions (linker 3 - 127; linker-4 - 131; linker-5-135) at the fourth and fifth positions in the 5'-flanking sequences to the CpG dinucleotide and the corresponding 5'-truncated versions 128,


132, and 136, respectively, were tested for their ability to induce cytokine secretion in BALB/c mouse spleen cell cultures. The levels of IL-12 and IL-6 secreted at 1.0 µg/mL concentration are shown in Figure 17. The results presented in Figure 17 suggest that the immunostimulatory activity is dependent on the nature of the linker incorporated. The substitution of the fourth and fifth nucleosides with C4-linker 3 (CpG DNA 127) had an insignificant effect on cytokine secretion compared with parent CpG DNA 4, suggesting that the nucleobase and sugar ring at these positions are not required for receptor recognition and/or binding. The deletion of the nucleotides beyond the linker substitutions (CpG DNA 128) caused higher IL-12 and IL-6 secretion than that found with CpG DNAs 4 and 127. As expected, the substitution of two C6-linkers (4) resulted in IL-12 secretion lower than and IL-6 secretion similar to that induced by parent CpG DNA 4. The 5'-truncated CpG DNA 132 induced higher cytokine secretion than did CpG DNA 131. The CpG DNAs 135 and 136, which had two C9-linkers (5), induced insignificant cytokine secretion, confirming the results obtained with mono-substituted CpG DNA containing the same linker as described above.

Example 14: Effect of Phosphodiester Linkages on Cytokine Induction

[0140] To test the effect of phosphodiester linkages on immunomer-induced cytokine induction, the following molecules were synthesized.

Table 21 PO-Immunomer sequences and analytical data

CpG DNA	Sequence ^a	Backbone ^b	Molecular Weight	
			Calculated	Found ^c
4	5'-CTATCTGACGTTCTCTGT-3'	PS	5702	5704
155	5'-CTATCTGACGTTCTCTGT-3'	PO	5432	5428
156	5'-CTGACGTTCTCTGT-X-TGTCTCTTGCAGTC-5'	PO	8656	8649
157	5'-YYCTGACGTTCTCTGT-X-TGTCTCTTGCAGTCYY-5'	PO	9208	9214



^aArrows indicate 5'-3' directionality of CpG dinucleotide in each DNA molecule and structures of X and Y are shown in boxes.

^bPS and PO stand for phosphorothioate and phosphodiester backbones, respectively.

^cAs determined by MALDI-TOF mass spectrometry.

[0141] PS-CpG DNA 4 (Table 21) was found to induce an immune response in mice (data not shown) with PO-CpG DNA 155 serving as a control. PO-immunomers 156 and 157 each contain two identical, truncated copies of the parent CpG DNA 155 joined through their 3'-ends via a glyceryl linker, X (Table 21). While 156 and 157 each contain the same oligonucleotide segments of 14 bases, the 5'-ends of 157 were modified by the addition of two C3-linkers, Y (Table 21). All oligonucleotides 4, 155-157 contain a 'GACGTT' hexameric motif known to activate the mouse immune system. [0142] The stability of PO-immunomers against nucleases was assessed by incubating CpG DNAs 4, 155-157 in cell culture medium containing 10% fetal bovine serum (FBS) (non-heat-inactivated) at 37 °C for 4, 24, and 48 hr. Intact CpG DNA remaining in the reaction mixtures were then determined by CGE. Figure 18 A-D shows the nuclease digestion profiles of CpG DNAs 4, 155-157 incubated in 10% FBS for 24 hr. The amount of full-length CpG DNA remaining at each time point is shown in Figure 18 E. As expected, the parent PS-CpG DNA 4 is the most resistant to serum nucleases. About 55% of 18-mer 4 remained undegraded after 48 hr incubation. In contrast, only about 5% of full-length PO-immunomer 155 remained after 4 hr under the same experimental conditions confirming that DNA containing phosphodiester linkages undergoes rapid degradation. As expected, both PO-immunomers 156 and 157 were more resistant than 155 to serum nucleases. After 4 hr, about 62% and 73% of 156 and 157 respectively were intact compared with about 5% of 155 (Fig. 18 E). Even after 48 hr, about 23% and 37% of 156 and 157, respectively, remained undegraded. As well as showing that 3'-3'-linked PO-immunomers are more stable against serum nucleases, these studies indicate that chemical modifications at the 5'-end can further increase nuclease stability.

[0143] The immunostimulatory activity of CpG DNAs was studied in BALB/c and C3H/HeJ mice spleen cell cultures by measuring levels of cytokines IL-12 and IL-6 secreted. All CpG DNAs induced a concentration-dependent cytokine secretion in BALB/c mouse spleen cell cultures (Fig. 19). At 3 µg/mL, PS-CpG DNA 4 induced 2656±256 and 12234±1180 pg/mL of IL-12 and IL-6 respectively. The parent PO-CpG DNA 155 did not raise cytokine levels above background except at a concentration of 10 µg/mL. This observation is consistent with the nuclease stability assay results. In contrast, PO-immunomers 156 and 157 induced both IL-12 and IL-6 secretion in BALB/c mouse spleen cell

cultures.

[0144] The results presented in Figure 19 show a clear distinction in cytokine induction profiles of PS- and PO-CpG DNAs. PO-immunomers **156** and **157** induced higher levels of IL-12 than did PS-CpG DNA **4** in BALB/c mouse spleen cell cultures (Fig. 19A). In contrast, at concentrations up to 3 µg/mL, they produced negligible amounts of IL-6 (Fig. 19B). Even at the highest concentration (10 µg/mL), PO-immunomer **156** induced significantly less IL-6 than did PS-CpG DNA **4**. The presence of C3 linkers at the 5'-terminus of PO-immunomer **157** resulted in slightly higher levels of IL-6 secretion compared with **156**. However, importantly, the levels of IL-6 produced by PO-immunomer **157** are much lower than those induced by PS CpG DNA **4**. The inset of Figure 19A shows the ratio of IL-12 to IL-6 secreted at 3 µg/mL concentration. In addition to increasing IL-12 secretion, PO-immunomers **156** and **157** induced higher levels of IFN-γ than did PS-CpG DNA **4** in BALB/c mouse spleen cell cultures (data not shown).

[0145] The different cytokine profiles induced by PO- and PS-CpG DNAs in BALB/c mouse spleen cell cultures prompted us to study the pattern of cytokine induction of CpG DNAs in C3H/HeJ mouse spleen cell cultures (an LPS lower-responsive strain). All three CpG DNAs tested in this assay induced concentration-dependent cytokine secretion (Fig. 20A and B). Since PO-CpG DNA **155** failed to induce cytokine secretion in BALB/c mouse spleen cell cultures, it was not further tested in C3H/HeJ spleen cell cultures. Both PO-immunomers **156** and **157** induced higher IL-12 production than did PS-CpG DNA **4** (Fig. 20A). However, at concentrations up to 3 µg/mL, neither induced IL-6 production. At the highest concentration tested (10 µg/mL), both induced significantly less IL-6 than did PS-CpG DNA **4** (Fig. 20B). The ratio of IL-12 to IL-6 secreted is calculated to distinguish cytokine secretion profiles of PS and PO CpG DNAs (Fig. 20A inset). In addition, the C3H/HeJ spleen cell culture results suggest that the responses observed with CpG DNAs are not due to LPS contamination.

[0146] PS-CpG DNAs have been shown to induce potent antitumor activity *in vivo*. Since PO-CpG DNAs exhibited greater nuclease stability and induced higher levels of IL-12 and IFN-γ secretion in *in vitro* assays, we were interested to see if these desirable properties of PO-immunomers improve the antitumor activity *in vivo*. We administered PO-immunomer **157** subcutaneously at a dose of 0.5 mg/kg every other day to nude mice bearing tumor xenografts of MCF-7 breast cancer cells that express wild-type p53, or DU-145 prostate cancer cells that express mutated p53. PO-immunomer **157** gave 57% growth inhibition of MCF-7 tumors on day 15 compared with the saline control (Fig. 21A). It also produced 52% growth inhibition of DU-145 tumors on day 34 (Fig. 21B). These antitumor studies suggest that PO-immunomers of the proposed design exhibit potent antitumor activity *in vivo*.

Example 22: Short immunomers

To test the effects of short immunomers on cytokine induction, the following immunomers were used. These results show that immunomers as short as 5 nucleotides per segment are effective in inducing cytokine production.

[0147]


Table 22. Immunomer Structure and Immunostimulatory Activity in BABL/C Mouse Spleen Cell Cultures

Oligo No.	Sequences and Modification (5'-3')	Oligo Length/ or Each Chain	IL-12 (pg/mL)	IL-6 (pg/mL)
			10 µg/mL	10 µg/mL
4	5'-CTATCTGACGTTCTCTGT-3'	18mer	2731	4547
25	5'-CTATCTGTCGTTCTCTGT-3'	18mer	795	789
158	5'-TCTGACGTTCT-3' 5'-TCTGACGTTCT-3' X ₁	11mer	3490	5319
159	5'-TCTGTCGTTCT-3' 5'-TCTGTCGTTCT-3' X ₁	11mer	3265	4625
160	5'-TCGTTG-3' 5'-TCGTTG-3' X ₁	6mer	2085	2961
161	5'-TCGTTG-3'XX 5'-TCGTTG-3'XX X ₁	6mer	3169	5194
162	5'-TCGTTG-3'X 5'-TCGTTG-3'X X ₁	6mer	1015	705
163	5'-TCGTT-3'X 5'-TCGTT-3'X X ₁	5mer	2623	3619
164	5'-ACGTTG-3'X 5'-ACGTTG-3'X X ₁	6mer	564	845
165	5'-GCGTTG-3'X 5'-GCGTTG-3'X X ₁	6mer	196	0
166	5'-CCGTTG-3'X 5'-CCGTTG-3'X X ₁	6mer	219	0
167	5'-GTCGTT-3'X 5'-GTCGTT-3'X X ₁	6mer	1441	5056
168	5'-TGTCGT-3'X 5'-TGTCGT-3'X X ₁	6mer	198	0
169	5'-TCGTTG-3'X 5'-TCGTTG-3'X X ₁ -X ₃ '-GTTGCT-5'	6mer	2410	4857

Normal phase represents a phosphorothioate linkage.


5

10



15

20



EQUIVALENTS

25

[0148] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention and appended claims.

30

35

40

45

50

55

SEQUENCE LISTING

5 <110> Hybridon, Inc.

<120> Modulation of Immunostimulatory Properties of
Oligonucleotide-Based Compounds By Optimal Presentation
of 5' Ends

10 <130> 351-10

<140> US 10/279,684
<141> 2002-10-24

<160> 115

15 <170> FastSEQ for Windows Version 4.0

<210> 1
<211> 16
<212> DNA
20 <213> Artificial Sequence

<220>
<223> immunomer sequence

25 <400> 1
gagaacgctc gacctt 16

<210> 2
<211> 16
<212> DNA
30 <213> Artificial Sequence

<220>
<223> immunomer sequence

35 <400> 2
ttccagctcg caagag 16

<210> 3
<211> 18
<212> DNA
40 <213> Artificial Sequence

<220>
<223> immunomer sequence

45 <400> 3
ctatctgacg ttctctgt 18

<210> 4
<211> 18
<212> DNA
50 <213> Artificial Sequence

<220>
<223> immunomer sequence

<221> modified_base
<222> 5
55 <223> C3-alkyl linker

	<400> 4	
	ctatntgacg ttctctgt	18
5	<210> 5	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> immunomer sequence	
	<221> modified_base	
	<222> 9	
	<223> 5OHdC	
15	<400> 5	
	ctatctgayg ttctctgt	18
	<210> 6	
	<211> 18	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
	<221> modified_base	
25	<222> 10	
	<223> 7-deaza-dG	
	<400> 6	
	ctatctgacr ttctctgt	18
30	<210> 7	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> immunomer sequence	
	<221> modified_base	
	<222> 4	
40	<223> C3-alkyl linker	
	<221> modified_base	
	<222> 9	
	<223> 5OHdC	
45	<400> 7	
	ctanctgayg ttctctgt	18
	<210> 8	
	<211> 18	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
	<221> modified_base	
55	<222> 4	
	<223> C3-alkyl linker	

5 <221> modified_base
 <222> 10
 <223> 7-deaza-dG

 <400> 8
 ctanctgacr ttctctgt 18

 10 <210> 9
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> immunomer sequence
 15
 <400> 9
 ctgacgttct ctgt 14

 <210> 10
 <211> 14
 20 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> immunomer sequence
 25
 <221> modified_base
 <222> 5
 <223> 5OHdC

 <400> 10
 30 ctgaygttct ctgt 14

 <210> 11
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> immunomer sequence

 <221> modified_base
 <222> 6
 40 <223> 7-deaza-dG

 <400> 11
 ctgacrttct ctgt 14

 45 <210> 12
 <211> 15
 <212> DNA
 <213> Artificial Sequence

 <220>
 50 <223> immunomer sequence

 <221> modified_base
 <222> (1)...(2)
 <223> 1',2'-dideoxyriboside

 55 <400> 12
 nntgacgttc tctgt 15

	<210> 13	
	<211> 16	
5	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
10	<221> modified_base	
	<222> (1)...(3)	
	<223> 1',2'-dideoxyriboside	
	<400> 13	
15	nnntgacgtt ctctgt	16
	<210> 14	
	<211> 16	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> immunomer sequence	
	<221> modified_base	
	<222> (1)...(3)	
25	<223> 1',2'-dideoxyriboside	
	<221> modified_base	
	<222> 7	
	<223> 5OHdC	
30	<400> 14	
	nnntgaygtt ctctgt	16
	<210> 15	
	<211> 16	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
40	<221> modified_base	
	<222> (1)...(3)	
	<223> 1',2'-dideoxyriboside	
	<221> modified_base	
	<222> 8	
45	<223> 7-deaza-dG	
	<400> 15	
	nnntgacrtt ctctgt	16
	<210> 16	
50	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
55	<400> 16	

tctgacgttc t 11

5 <210> 17
 <211> 14
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> immunomer sequence

<221> modified_base
 <222> (1)...(3)
 <223> 1',2'-dideoxyriboside

15 <400> 17
 nnntctgacg ttct 14

<210> 18
 <211> 14
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> immunomer sequence

25 <221> modified_base
 <222> (1)...(3)
 <223> 1',2'-dideoxyriboside

<221> modified_base
 <222> 9
 <223> 5OHdC

30 <400> 18
 nnntctgayg ttct 14

<210> 19
 <211> 14
 <212> DNA
 <213> Artificial Sequence

35 <220>
 <223> immunomer sequence

40 <221> modified_base
 <222> (1)...(3)
 <223> 1',2'-dideoxyriboside

45 <221> modified_base
 <222> 10
 <223> 7-deaza-dG

<400> 19
 nnntctgacr ttct 14

50 <210> 20
 <211> 18
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> immunomer sequence

	<400> 20	
	ctatctgtcg ttctctgt	18
5	<210> 21	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> immunomer sequence	
	<221> modified_base	
	<222> 7	
	<223> 7-deazaguanosine	
15	<400> 21	
	tctgtcrttc t	11
	<210> 22	
	<211> 11	
20	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
25	<221> modified_base	
	<222> 7	
	<223> deoxyarabinoguanosine	
	<400> 22	
30	tctgtcrttc t	11
	<210> 23	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> immunomer sequence	
	<221> modified_base	
	<222> 7	
40	<223> arabinocytosine	
	<221> modified_base	
	<222> 8	
	<223> deoxyarabinoguanosine	
45	<400> 23	
	tctgtyrttc t	11
	<210> 24	
	<211> 13	
50	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
55	<221> modified_base	
	<222> (1)...(2)	
	<223> propoxyphosphorothioate	

5	<221> modified_base <222> 9 <223> 7-deazaguanosine	
	<400> 24 nntctgtcrt tct	13
10	<210> 25 <211> 14 <212> DNA <213> Artificial Sequence	
15	<220> <223> immunomer sequence	
	<221> modified_base <222> 6 <223> dexoyarabinoguanosine	
20	<400> 25 ctgtcrttct ctgt	14
25	<210> 26 <211> 14 <212> DNA <213> Artificial Sequence	
	<220> <223> immunomer sequence	
30	<221> modified_base <222> 5 <223> arabinocytosine	
35	<221> modified_base <222> 6 <223> deoxyarabinoguanosine	
	<400> 26 ctgtyrttct ctgt	14
40	<210> 27 <211> 13 <212> DNA <213> Artificial Sequence	
45	<220> <223> immunomer sequence	
	<221> modified_base <222> (1)...(2) <223> propoxyphosphorothioate	
50	<221> modified_base <222> 9 <223> 7-deazaguanosine	
	<400> 27 nntctgacrt tct	13
55	<210> 28 <211> 11	

	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> immunomer sequence	
	<221> modified_base	
	<222> 7	
	<223> deoxyarabinoguanosine	
10	<400> 28	
	tctgacrttc t	11
	<210> 29	
	<211> 11	
15	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
20	<221> modified_base	
	<222> 6	
	<223> arabinocytosine	
	<221> modified_base	
25	<222> 7	
	<223> deoxyarabinoguanosine	
	<400> 29	
	tctgayrttc t	11
30	<210> 30	
	<211> 14	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
35	<223> immunomer sequence	
	<221> modified_base	
	<222> 5	
	<223> arabinocytosine	
40	<400> 30	
	ctgaygttct ctgt	14
	<210> 31	
	<211> 14	
45	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
50	<221> modified_base	
	<222> 6	
	<223> deoxyarabinoguanosine	
	<400> 31	
55	ctgacrttct ctgt	14
	<210> 32	

<211> 14
 <212> DNA
 <213> Artificial Sequence
 5 <220>
 <223> immunomer sequence
 <221> modified_base
 <222> 5
 10 <223> arabinocytosine
 <221> modified_base
 <222> 6
 <223> deoxyarabinoguanosine
 15 <400> 32
 ctgayrttct ctgt 14
 <210> 33
 <211> 14
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> immunomer sequence
 25 <400> 33
 ctgtcgttct ctgt 14
 <210> 34
 <211> 12
 <212> DNA
 30 <213> Artificial Sequence
 <220>
 <223> immunomer sequence
 35 <400> 34
 ctgtcgttct ct 12
 <210> 35
 <211> 11
 <212> DNA
 40 <213> Artificial Sequence
 <220>
 <223> immunomer sequence
 45 <400> 35
 tctgtcgttc t 11
 <210> 36
 <211> 18
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <223> immunomer sequence
 55 <400> 36
 ctcactttcg ttctctgt 18
 <210> 37

<211> 14
 <212> DNA
 <213> Artificial Sequence
 5 <220>
 <223> immunomer sequence
 <400> 37
 ctttcgttct ctgt 14
 10 <210> 38
 <211> 12
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <223> immunomer sequence
 <400> 38
 ctttcgttct ct 12
 20 <210> 39
 <211> 11
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> immunomer sequence
 <221> modified_base
 <222> 6
 <223> 5-hydroxycytosine
 30 <400> 39
 tctttygttc t 11
 <210> 40
 <211> 11
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> immunomer sequence
 40 <221> modified_base
 <222> 7
 <223> 7-deazaguanosine
 <400> 40
 45 tctttcrttc t 11
 <210> 41
 <211> 11
 <212> DNA
 <213> Artificial Sequence
 50 <220>
 <223> immunomer sequence
 <221> modified_base
 <222> 6
 55 <223> 5-hydroxycytosine

	<400> 41	
	tctgtygttc t	11
5	<210> 42	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> immunomer sequence	
	<221> modified_base	
	<222> 7	
	<223> 7-deazaguanosine	
15	<400> 42	
	tctgtcrttc t	11
	<210> 43	
	<211> 18	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
	<221> modified_base	
25	<222> 9	
	<223> 5-hydroxycytosine	
	<400> 43	
	ctatctgayg ttctctgt	18
30	<210> 44	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> immunomer sequence	
	<221> modified_base	
	<222> 10	
40	<223> 7-deazaguanosine	
	<400> 44	
	ctatctgacr ttctctgt	18
	<210> 45	
45	<211> 14	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> immunomer sequence	
50	<221> modified_base	
	<222> 5	
	<223> 5-hydroxycytosine	
55	<400> 45	
	ctgaygttct ctgt	14

5	<210> 46 <211> 14 <212> DNA <213> Artificial Sequence	
	<220> <223> immunomer sequence	
10	<221> modified_base <222> 6 <223> 7-deazaguanosine	
	<400> 46 ctgacrttct ctgt	14
15	<210> 47 <211> 11 <212> DNA <213> Artificial Sequence	
20	<220> <223> immunomer sequence	
	<221> modified_base <222> 6 <223> 5-hydroxycytosine	
25	<400> 47 tctgaygttc t	11
30	<210> 48 <211> 11 <212> DNA <213> Artificial Sequence	
	<220> <223> immunomer sequence	
35	<221> modified_base <222> 7 <223> 7-deazaguanosine	
40	<400> 48 tctgacrttc t	11
	<210> 49 <211> 10 <212> DNA <213> Artificial Sequence	
45	<220> <223> immunomer sequence	
	<400> 49 ctgacgttct	10
50	<210> 50 <211> 16 <212> DNA <213> Artificial Sequence	
55	<220> <223> immunomer sequence	

5
 <221> modified_base
 <222> (1)...(2)
 <223> C3-linker

<400> 50
 nnctgacgtt ctctgt 16

10
 <210> 51
 <211> 11
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotides in 10% Bovine Serum PBS solution

15
 <221> modified_base
 <222> 7
 <223> dGdeaza

20
 <400> 51
 tctgtcgttc t 11

<210> 52
 <211> 16
 <212> DNA
 <213> Artificial Sequence

25
 <220>
 <223> oligonucleotides in 10% Bovine Serum PBS solution

<221> modified_base
 <222> (1)...(2)
 <223> C3-linker

30
 <400> 52
 nnctgtcgtt ctctgt 16

35
 <210> 53
 <211> 15
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotides in 10% Bovine Serum PBS solution

40
 <221> modified_base
 <222> 1
 <223> uridine

45
 <221> modified_base
 <222> 2
 <223> 2'-OMe-ribonucleoside

<400> 53
 uctgtcgttc tctgt 15

50
 <210> 54
 <211> 19
 <212> DNA
 <213> Artificial Sequence

55
 <220>

	<223> phosphorothioate CpG DNA sequence	
	<400> 54	
5	tccatgacgt tcctgatgc	19
	<210> 55	
	<211> 17	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> phosphorothioate CpG DNA sequence	
	<400> 55	
15	ccatgacgtt cctgatg	17
	<210> 56	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> phosphorothioate CpG DNA sequence	
	<400> 56	
25	tccatgacgt tcctgatg	18
	<210> 57	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> phosphorothioate CpG DNA sequence	
	<400> 57	
35	ccatgacgtt cctgatgc	18
	<210> 58	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> CpG DNA sequence	
	<400> 58	
	cctactagcg ttctcatc	18
45	<210> 59	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> CpG DNA sequence	
	<221> misc_feature	
	<222> 10	
	<223> alkyl linker 2 of Figure 14	
55	<400> 59	
	cctactagcn ttctcatc	18

```

5      <210> 60
      <211> 18
      <212> DNA
      <213> Artificial Sequence

      <220>
      <223> CpG DNA sequence

10     <221> misc_feature
      <222> 7
      <223> alkyl linker 2 of Figure 14

      <400> 60
      cctactngcg ttctcatc 18

15     <210> 61
      <211> 18
      <212> DNA
      <213> Artificial Sequence

20     <220>
      <223> CpG DNA sequence

      <221> misc_feature
      <222> 5
      <223> alkyl linker 2 of Figure 14

25     <400> 61
      cctantagcg ttctcatc 18

30     <210> 62
      <211> 18
      <212> DNA
      <213> Artificial Sequence

      <220>
      <223> CpG DNA sequence

35     <221> misc_feature
      <222> (4)...(5)
      <223> alkyl linker 2 of Figure 14

      <400> 62
      cctnntagcg ttctcatc 18

40     <210> 63
      <211> 18
      <212> DNA
      <213> Artificial Sequence

45     <220>
      <223> CpG DNA sequence

      <221> misc_feature
      <222> (1)...(2)
      <223> alkyl linker 2 of Figure 14

      <400> 63
      nntactagcg ttctcatc 18

55     <210> 64
      <211> 18

```

<212> DNA
 <213> Artificial Sequence

5 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 <222> 12
 <223> alkyl linker 2 of Figure 14

10 <400> 64
 cctactagcg tntcatc 18

 <210> 65
 <211> 18
 15 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

20 <221> misc_feature
 <222> 14
 <223> alkyl linker 2 of Figure 14

 <400> 65
 25 cctactagcg ttcncatc 18

 <210> 66
 <211> 18
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 <222> (14)...(15)
 35 <223> alkyl linker 2 of Figure 14

 <400> 66
 cctactagcg ttcnnatc 18

 <210> 67
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 45 <223> CpG DNA sequence

 <221> misc_feature
 <222> 4
 <223> ethylene-glycol linker 6 of Figure 14

50 <400> 67
 cctnctagcg ttctcatc 18

 <210> 68
 <211> 18
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> CpG DNA sequence
 5 <221> misc_feature
 <222> 14
 <223> ethylene-glycol linker 6 of Figure 14
 <400> 68
 cctactagcg ttcncatc 18
 10 <210> 69
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 4
 <223> ethylene-glycol linker 7 of Figure 14
 20 <400> 69
 cctnctagcg ttctcatc 18
 <210> 70
 <211> 19
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 30 <221> misc_feature
 <222> 14
 <223> ethylene-glycol linker 7 of Figure 14
 <400> 70
 35 cctactagcg ttctncatc 19
 <210> 71
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 5
 45 <223> alkyl linker 1 of Figure 14
 <400> 71
 ctatntgacg ttctctgt 18
 50 <210> 72
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 55 <223> CpG DNA sequence

<221> misc_feature
 <222> 4
 <223> alkyl linker 1 of Figure 14
 5
 <400> 72
 ctanctgacg ttctctgt 18
 <210> 73
 <211> 18
 10 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 15 <221> misc_feature
 <222> 8
 <223> alkyl linker 2 of Figure 14
 <400> 73
 20 ctatctgncg ttctctgt 18
 <210> 74
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 6
 30 <223> alkyl linker 2 of Figure 14
 <400> 74
 ctatcngacg ttctctgt 18
 <210> 75
 35 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 40 <223> CpG DNA sequence
 <221> misc_feature
 <222> 4
 <223> alkyl linker 2 of Figure 14
 45 <400> 75
 ctanctgacg ttctctgt 18
 <210> 76
 <211> 18
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 55 <222> (1)...(5)
 <223> alkyl linker 2 of Figure 14

	<400> 76 nnnnntgacg ttctctgt	18
5	<210> 77 <211> 17 <212> DNA <213> Artificial Sequence	
10	<220> <223> CpG DNA sequence	
	<221> misc_feature <222> (1)...(4) <223> alkyl linker 2 of Figure 14	
15	<400> 77 nnnntgacgt tctctgt	17
	<210> 78 <211> 16 <212> DNA <213> Artificial Sequence	
20	<220> <223> CpG DNA sequence	
25	<221> misc_feature <222> (1)...(3) <223> alkyl linker 2 of Figure 14	
30	<400> 78 nnntgacgtt ctctgt	16
	<210> 79 <211> 15 <212> DNA <213> Artificial Sequence	
35	<220> <223> CpG DNA sequence	
	<221> misc_feature <222> (1)...(2) <223> alkyl linker 2 of Figure 14	
40	<400> 79 nntgacgttc tctgt	15
45	<210> 80 <211> 14 <212> DNA <213> Artificial Sequence	
50	<220> <223> CpG DNA sequence	
	<221> misc_feature <222> 1 <223> alkyl linker 2 of Figure 14	
55	<400> 80 ntgacgttct ctgt	14

5 <210> 81
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

 10 <221> misc_feature
 <222> 5
 <223> alkyl linker 3 of Figure 14

 <400> 81
 15 ctatntgacg ttctctgt 18

 <210> 82
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 20 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 <222> 4
 25 <223> alkyl linker 3 of Figure 14

 <400> 82
 ctanctgacg ttctctgt 18

 <210> 83
 30 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence
 35 <221> misc_feature
 <222> (4)...(5)
 <223> alkyl linker 3 of Figure 14

 <400> 83
 40 ctanntgacg ttctctgt 18

 <210> 84
 <211> 15
 <212> DNA
 45 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 50 <222> (1)...(2)
 <223> alkyl linker 3 of Figure 14

 <400> 84
 nntgacgttc tctgt 15

 55 <210> 85
 <211> 18

<212> DNA
 <213> Artificial Sequence

 5 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 <222> 5
 <223> alkyl linker 4 of Figure 14

 10 <400> 85
 ctatntgacg ttctctgt 18

 <210> 86
 <211> 18
 15 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

 20 <221> misc_feature
 <222> 4
 <223> alkyl linker 4 of Figure 14

 <400> 86
 25 ctanctgacg ttctctgt 18

 <210> 87
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 30 <220>
 <223> CpG DNA sequence

 <221> misc_difference
 <222> (4)...(5)
 35 <223> alkyl linker 4 of Figure 14

 <400> 87
 ctanntgacg ttctctgt 18

 <210> 88
 40 <211> 15
 <212> DNA
 <213> Artificial Sequence

 <220>
 45 <223> CpG DNA sequence

 <221> misc_feature
 <222> (1)...(2)
 <223> alkyl linker 4 of Figure 14

 50 <400> 88
 nntgacgttc tctgt 15

 <210> 89
 <211> 18
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> CpG DNA sequence
 5 <221> misc_feature
 <222> 5
 <223> alkyl linker 5 of Figure 14
 <400> 89
 ctatntgacg ttctctgt 18
 10 <210> 90
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 4
 20 <223> alkyl linker 5 of Figure 14
 <400> 90
 ctanctgacg ttctctgt 18
 <210> 91
 <211> 18
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 30 <221> misc_feature
 <222> (4)...(5)
 <223> alkyl linker 5 of Figure 14
 <400> 91
 35 ctanntgacg ttctctgt 18
 <210> 92
 <211> 15
 <212> DNA
 40 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 45 <222> (1)...(2)
 <223> alkyl linker 5 of Figure 14
 <400> 92
 nntgacgttc tctgt 15
 50 <210> 93
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> CpG DNA sequence

<221> misc_feature
 <222> 4
 <223> ethylene-glycol linker 6 of Figure 14
 5 <400> 93
 ctanctgacg ttctctgt 18
 <210> 94
 <211> 18
 10 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 15 <221> misc_difference
 <222> 14
 <223> ethylene-glycol linker 6 of Figure 14
 <400> 94
 ctatctgacg ttcnctgt 18
 20 <210> 95
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 4
 30 <223> ethylene-glycol linker 7 of Figure 14
 <400> 95
 ctanctgacg ttctctgt 18
 <210> 96
 35 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 40 <221> misc_feature
 <222> 14
 <223> ethylene-glycol linker 7 of Figure 14
 <400> 96
 45 ctatctgacg ttcnctgt 18
 <210> 97
 <211> 18
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 <221> misc_difference
 55 <222> 8
 <223> branched alkyl linker 8 of Figure 14

	<400> 97	
	ctatctgncg ttctctgt	18
5	<210> 98	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> CpG DNA sequence	
	<221> misc_feature	
	<222> 7	
	<223> branched alkyl linker 8 of Figure 14	
15	<400> 98	
	ctatctnacg ttctctgt	18
	<210> 99	
	<211> 18	
20	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> CpG DNA sequence	
25	<221> misc_feature	
	<222> 6	
	<223> branched alkyl linker 8 of Figure 14	
30	<400> 99	
	ctatcngacg ttctctgt	18
	<210> 100	
	<211> 18	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
	<223> CpG DNA sequence	
	<221> misc_feature	
40	<222> 5	
	<223> branched alkyl linker 8 of Figure 14	
	<400> 100	
	ctatntgacg ttctctgt	18
45	<210> 101	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> CpG DNA sequence	
	<221> misc_feature	
	<222> 4	
	<223> branched alkyl linker 8 of Figure 14	
55	<400> 101	
	ctanctgacg ttctctgt	18

5 <210> 102
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

 10 <221> misc_feature
 <222> 11
 <223> branched alkyl linker 8 of Figure 14

 <400> 102
 ctatctgacg ntctctgt 18
 15
 <210> 103
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 20 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 <222> 12
 <223> branched alkyl linker 8 of Figure 14
 25
 <400> 103
 ctatctgacg tntctctgt 18

 <210> 104
 30 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence
 35
 <221> misc_feature
 <222> 13
 <223> branched alkyl linker 8 of Figure 14

 <400> 104
 40 ctatctgacg ttntctgt 18

 <210> 105
 <211> 18
 <212> DNA
 45 <213> Artificial Sequence

 <220>
 <223> CpG DNA sequence

 <221> misc_feature
 50 <222> 14
 <223> branched alkyl linker 8 of Figure 14

 <400> 105
 ctatctgacg ttcnctgt 18
 55
 <210> 106
 <211> 18

<212> DNA
 <213> Artificial Sequence
 5 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 8
 <223> branched alkyl linker 9 of Figure 14
 10 <400> 106
 ctatctgncg ttctctgt 18
 <210> 107
 <211> 18
 15 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> CpG DNA sequence
 20 <221> misc_feature
 <222> 7
 <223> branched alkyl linker 9 of Figure 14
 <400> 107
 25 ctatctnacg ttctctgt 18
 <210> 108
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <223> CpG DNA sequence
 <221> misc_feature
 <222> 4
 35 <223> branched alkyl linker 9 of Figure 14
 <400> 108
 ctanctgacg ttctctgt 18
 <210> 109
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> CpG DNA sequence
 <221> misc_difference
 <222> 12
 <223> branched alkyl linker 9 of Figure 14
 50 <400> 109
 ctatctgacg tntctgt 18
 <210> 110
 <211> 18
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> CpG DNA sequence
 5 <221> misc_feature
 <222> 14
 <223> branched alkyl linker 9 of Figure 14
 <400> 110
 ctatctgacg ttcnctgt 18
 10 <210> 111
 <211> 29
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <223> immunomer sequence
 <221> modified_base
 <222> 15
 <223> glyceryldiphosphodiester
 20 <400> 111
 ctgacgttct ctgtntgtct cttgcagtc 29
 <210> 112
 <211> 33
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> immunomer sequence
 30 <221> modified_base
 <222> 1,2,32,33
 <223> propoxyphosphodiester
 <221> modified_base
 35 <222> 17
 <223> glyceryldiphosphodiester
 <400> 112
 yyctgacgtt ctctgtntgt ctcttgcaagt cyy 33
 40 <210> 113
 <211> 11
 <212> DNA
 <213> Artificial Sequence
 45 <220>
 <223> immunomer sequence
 <400> 113
 tctgacgttc t 11
 50 <210> 114
 <211> 11
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> immunomer sequence

<400> 114
tctgtcgttc t

11

5 <210> 115
<211> 18
<212> DNA
<213> Artificial Sequence

10 <220>
<223> immunomer sequence


<400> 115
tgtctcttgc agtctatc

18

15

Claims

- 20 1. An immunomer, comprising at least two oligonucleotides linked together by a non-nucleotidic linker, wherein at least one of the oligonucleotides is an immunostimulatory oligonucleotide having an accessible 5' end and comprising an immunostimulatory dinucleotide.
- 25 2. The immunomer according to claim 1 wherein the immunostimulatory dinucleotide is selected from the group consisting of CpG, C*pG, CpG*, and C*pG*, wherein C is cytidine or 2'-deoxycytidine, C* is 2'-deoxythymidine, arabinocytidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside, G is guanosine or 2'-deoxyguanosine, G* is 2' deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, or other non-natural purine nucleoside, and p is an internucleoside linkage selected from the group consisting of phosphodiester, phosphorothioate, and phosphorodithioate.
- 30 3. The immunomer of claim 1, wherein the immunostimulatory oligonucleotide comprises an immunostimulatory domain having the structure
- 35



wherein:

40

Y is cytidine, 2' deoxycytidine arabinocytidine, 2'-deoxythymidine, 2'-deoxy-2'-substitutedarabinocytidine, 2'-O-substitutedarabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside;

45

Z is guanosine or 2'-deoxyguanosine, G* is 2' deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted-arabinoguanosine, 2'-O-substituted-arabinoguanosine, 2'-deoxyinosine or other non-natural purine nucleoside,

50

N1, at each occurrence, is preferably a naturally occurring or a synthetic nucleoside or an immunostimulatory moiety;


Nn, at each occurrence, is a naturally occurring nucleoside or an immunostimulatory moiety;

55


provided that at least one N1 or Nn is an immunostimulatory moiety and that the 5' N1 includes a nucleobase; and

wherein n is a number from 0-30.

4. The immunomer according to claim 3, wherein the immunostimulatory moiety is selected from the group consisting of abasic nucleosides, arabinonucleosides, 2'-deoxyuridine, α -deoxyribonucleosides, β -L-deoxyribonucleosides, and nucleosides linked by a modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleotide linkage being selected from the group consisting of C2-C18 alkyl linker, poly(ethylene glycol) linker, 2-aminobutyl-1,3-propanediol linker, 2'-5' internucleoside linkage, methylphosphonate internucleoside linkage; methylphosphonothioates, phosphotriesters, phosphothiotriesters, phosphorothioates, phosphorodithioates, triester prodrugs, sulfones, sulfonamides, sulfamates, formacetal, N-methylhydroxylamine, carbonate, carbamate, morpholino, boranophosphonate, phosphoramidates, especially primary amino-phosphoramidates, N3 phosphoramidates and N5 phosphoramidates, and stereospecific linkages, nucleosides having sugar modifications, 2'-substituted pentose sugars including, without limitation, 2'-O-methylribose, 2'-O-methoxyethylribose, 2'-O-propargylribose, and 2'-fluororibose; 3'-substituted pentose sugars, including, without limitation, 3'-O-methylribose; 1',2'-dideoxyribose; arabinose; substituted arabinose sugars, hexose sugars, and alpha-anomers, peptide nucleic acids (PNA), peptide nucleic acids with phosphate groups (PHONA), locked nucleic acids (LNA), morpholinonucleic acids, and oligonucleotides having backbone linker sections having a length of from about 2 angstroms to about 200 angstroms, alkyl linkers or amino linkers, DNA isoforms, β -L-deoxyribonucleosides, α -deoxyribonucleosides, nucleosides having unnatural internucleoside linkage positions, and nucleosides having modified heterocyclic bases.
5. The immunomer of claim 1, wherein each of the oligonucleotides is independently is linked to a non-nucleotidic linker at the 3'end of the oligonucleotide, at an internucleotide linkage of the oligonucleotide, at a functionalized nucleobase of the oligonucleotide or at a functionalized sugar of the oligonucleotide.
6. The immunomer according to claim 1 having the structure



7. The immunomer of claim 1 wherein the immunomer comprises at least one oligonucleotide that is complementary to a gene.
8. The immunomer of claim 1 wherein the immunomer comprises at least one ribozyme or a decoy oligonucleotide.
9. The immunomer of claim 3 wherein at least one Nn includes a G4 tetranucleotide.
10. The immunomer of claim 2 or 3 wherein the non-naturally occurring pyrimidine has the structure (I):



(I)

wherein:

D is a hydrogen bond donor;

D' is selected from the group consisting of hydrogen, hydrogen bond donor, hydrogen bond acceptor, hydrophilic group, hydrophobic group, electron withdrawing group and electron donating group, excluding bromine; A is a hydrogen bond acceptor or a hydrophilic group;

A is a hydrogen bond acceptor or a hydrophilic group;

A' is selected from the group consisting of hydrogen bond acceptor, hydrophilic group, hydrophobic group, electron withdrawing group and electron donating group;

X is carbon or nitrogen; and

S' is a pentose or hexose sugar ring or a non-naturally occurring sugar.

11. The immunomer according to claim 10 wherein the sugar ring is derivatized with a phosphate moiety, modified phosphate moiety, or other non-nucleotidic linker moiety suitable for linking the pyrimidine nucleoside to another nucleoside or nucleoside analog.


12. The immunomer according to claim 10 wherein the hydrogen bond donors are selected from the group consisting of -NH-, -NH₂, -SH and -OH.

13. The immunomer according to claim 10 wherein the hydrogen bond acceptors are selected from the group consisting of C=O, C=S, and the ring nitrogen atoms of an aromatic heterocycle.

14. The immunomer according to claim 10 wherein the non-naturally occurring pyrimidine base is selected from the group consisting of 5-hydroxycytosine, 5-hydroxymethylcytosine, N4-alkylcytosine, N4-ethylcytosine, and 4-thiouracil.

15. The immunomer according to claim 10 wherein the non-naturally occurring sugar is selected from arabinose and arabinose analogs.

16. The immunomer according to claim 2 or 3 wherein the non-naturally occurring purine nucleoside has the structure (II):



(II)

wherein:

D is a hydrogen bond donor;

D' is selected from the group consisting of hydrogen, hydrogen bond donor, and hydrophilic group;

A is a hydrogen bond acceptor or a hydrophilic group;

X is carbon or nitrogen;

each L is independently selected from the group consisting of C, O, N and S; and

S' is a pentose or hexose sugar ring, or a non-naturally occurring sugar.

17. The immunomer according to claim 16 wherein the sugar ring is derivatized with a phosphate moiety, modified phosphate moiety, or other linker moiety suitable for linking the pyrimidine nucleoside to another nucleoside or nucleoside analog.

18. The immunomer according to claim 16 wherein the hydrogen bond donors are selected from the group consisting of -NH-, -NH₂, -SH and -OH.

19. The immunomer according to claim 16 wherein the hydrogen bond acceptors are selected from the group consisting of C=O, C=S, -N= and the ring nitrogen atoms of an aromatic heterocycle.

20. The immunomer according to claim 16 wherein the non-naturally occurring purine is 6-thioguanine or 7-deazaguanine.

21. The immunomer according to claim 1, wherein the non-nucleotidic linker is selected from the group consisting of a linker from about 2 angstroms to about 200 angstroms in length, a metal, a soluble or insoluble biodegradable polymer bead, an organic moiety having functional groups that permit attachment to the 3'-terminal nucleoside of the oligonucleotide, a biomolecule, a cyclic or acyclic small molecule, an aliphatic or aromatic hydrocarbon, either of which optionally can include, either in the linear chain connecting the oligonucleotides or appended to it, one or more functional groups selected from the group consisting of hydroxy, amino, thiol, thioether, ether, amide, thioamide, ester, urea, and thiourea; amino acids, carbohydrates, cyclodextrins, adamantane, cholesterol, haptens, antibiotics, glycerol or a glycerol homolog of the formula HO-(CH₂)_o-CH(OH)-(CH₂)_p-OH, wherein o and p independently are integers from 1 to about 6, and a derivative of 1,3-diamino-2-hydroxypropane.

22. The immunomer according to claim 2 or 3, wherein the internucleoside linkages consist essentially of phosphodiester linkages.

23. An immunomer conjugate, comprising an immunomer, according to claim 1 and an antigen conjugated to the immunomer at a position other than the accessible 5' end.

24. The immunomer according to claim 2, wherein C*G* is arabinocytosine or 2'-deoxy-2-substituted arabinocytosine and arabinoguanosine or 2'-deoxy-2'-substituted arabinoguanosine, 2'-deoxy-7-deazaguanosine or 2'-deoxy-6-thi-


oguanosine, or 2'-deoxyinosine.

25. A pharmaceutical formulation comprising an immunomer according to claim 1 and a physiologically acceptable carrier.
26. A method for generating an immune response in a vertebrate, the method comprising administering to the vertebrate an immunomer according to claim 1.
27. A method for generating an immune response in a vertebrate, the method comprising administering to the vertebrate an immunomer conjugate according to claim 23.
28. A method for therapeutically treating a patient having a disease or disorder, such method comprising administering to the patient an immunomer according to claim 1.
29. The method according to claim 28 wherein the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, skin disorders, allergy, asthma or a disease caused by a pathogen.
30. A method for therapeutically treating a patient having a disease or disorder, such method comprising administering to the patient an immunomer conjugate according to claim 23.
31. A method for therapeutically treating a patient having a disease or disorder, such method comprising administering to the patient an immunomer according to claim 6.
32. The method according to claim 30 wherein the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, allergy, asthma or a disease caused by a pathogen.
33. The method according to claim 31 wherein the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, allergy, asthma or a disease caused by a pathogen.
34. The method of claim 26 further comprising administering a vaccine.
35. The method of claim 26, wherein the immunomer or the vaccine, or both, are linked to an immunogenic protein.
36. The method of claim 26 further comprising administering an adjuvant.
37. An immunomer, comprising at least two oligonucleotides linked directly together, wherein at least one of the oligonucleotides is an immunostimulatory oligonucleotide having an accessible 5' end and comprising an immunostimulatory dinucleotide, and wherein at least one of the oligonucleotides comprises at least one non-naturally occurring purine or at least one non-naturally occurring pyrimidine
38. The immunomer of claim 37 wherein each of said oligonucleotides of said immunomer is directly linked at the 3' end of the respective oligonucleotide, at an internucleotide linkage of the respective oligonucleotide, at a functionalized nucleobase of the respective oligonucleotide or at a functionalized sugar of the respective oligonucleotide.

Amended claims in accordance with Rule 86(2) EPC.

1. An immunomer, comprising at least two oligonucleotides linked together by a non-nucleotidic linker, wherein at least one of the oligonucleotides is an immunostimulatory oligonucleotide having an accessible 5' end and comprising an immunostimulatory dinucleotide.
2. The immunomer according to claim 1 wherein the immunostimulatory dinucleotide is selected from the group consisting of CpG, C*pG, CpG*, and C*pG*, wherein C is cytidine or 2'-deoxycytidine, C* is 2'-deoxythymidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, G is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, and p is an internucleoside linkage selected from the group consisting of phosphodiester, phosphorothioate, and phosphorodithioate.
3. The immunomer of claim 1, wherein the immunostimulatory oligonucleotide comprises an immunostimulatory

domain having the structure



wherein:

Y is cytidine, 2'-deoxycytidine arabinocytidine, 2'-deoxythymidine, 2'-deoxy-2'-substituted arabinocytidine, 2'-O-substituted arabinocytidine, 2'-deoxy-5-hydroxycytidine, 2'-deoxy-N4-alkyl-cytidine, 2'-deoxy-4-thiouridine or other non-natural pyrimidine nucleoside;

Z is guanosine or 2'-deoxyguanosine, G* is 2'-deoxy-7-deazaguanosine, 2'-deoxy-6-thioguanosine, arabinoguanosine, 2'-deoxy-2'-substituted arabinoguanosine, 2'-O-substituted arabinoguanosine, 2'-deoxyinosine or other non-natural purine nucleoside,

N1, at each occurrence, is preferably a naturally occurring or a synthetic nucleoside or an immunostimulatory moiety;

Nn, at each occurrence, is a naturally occurring nucleoside or an immunostimulatory moiety;

provided that at least one N1 or Nn is an immunostimulatory moiety and that the 5' N1 includes a nucleobase; and wherein n is a number from 0-30.

4. The immunomer according to claim 3, wherein the immunostimulatory moiety is selected from the group consisting of abasic nucleosides and nucleosides linked by a modified internucleoside linkage to the adjacent nucleoside on the 3' side, the modified internucleotide linkage being selected from the group consisting of C2-C18 alkyl linker, poly(ethylene glycol) linkage and 2-aminobutyl-1,3-propanediol linker.

5. The immunomer of claim 1, wherein each of the oligonucleotides is independently is linked to a non-nucleotidic linker.

6. The immunomer according to claim 5 wherein the sugar ring is derivatized with a non-nucleotidic linker moiety suitable for linking the pyrimidine nucleoside to another nucleoside or nucleoside analog.

7. The immunomer according to claim 1, wherein the non-nucleotidic linker is selected from the group consisting of an aliphatic or aromatic hydrocarbon, either of which optionally can include, either in the linear chain connecting the oligonucleotides or appended to it, one or more functional groups selected from the group consisting of hydroxy, amino, thiol, thioether, glycerol or a glycerol homolog of the formula $HO-(CH_2)_o-CH(OH)-(CH_2)_p-OH$, wherein o and p independently are integers from 1 to about 6, and a derivative of 1,3-diamino-2-hydroxypropane.

8. The immunomer according to claim 2 or 3, wherein the internucleoside linkages consist essentially of phosphodiester linkages.

9. An immunomer conjugate, comprising an immunomer, according to claim 1 and an antigen conjugated to the immunomer at a position other than the accessible 5' end.

10. A pharmaceutical formulation comprising an immunomer according to claim 1 and a physiologically acceptable carrier.

11. A method for generating an immune response in a vertebrate, the method comprising administering to the vertebrate an immunomer according to claim 1.

12. A method for therapeutically treating a patient having a disease or disorder, such method comprising administering to the patient an immunomer according to claim 1.

13. The method according to claim 12 wherein the disease or disorder to be treated is cancer, an autoimmune disorder, airway inflammation, inflammatory disorders, skin disorders, allergy, asthma or a disease caused by a pathogen.

14. An immunomer, comprising at least two oligonucleotides linked directly together, wherein at least one of the oligonucleotides is an immunostimulatory oligonucleotide having an accessible 5' end and comprising an immunostimulatory dinucleotide, and wherein at least one of the oligonucleotides comprises at least one non-naturally occurring purine or at least one non-naturally occurring pyrimidine

5

10

15

20

25

30

35

40

45

50

55

Figure 1

5'~~~~~3' Immunostimulatory oligonucleotide

Immunomers




Figure 2




Figure 3

Linkers for linear synthesis




Figure 4




Figure 5

Linear Synthesis of Immunomers




Figure 6

Parallel Synthesis of Immunomers




Figure 7A




Figure 7B




Figure 7C




Figure 8A




Figure 8B




Figure 9A




Figure 9B




Figure 9C




Figure 10A




Figure 10B




Figure 10C




Figure 11A




Figure 11B




Figure 11C




Figure 12





Figure 13






Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 03 01 7211

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 02 26757 A (HYBRIDON INC) 4 April 2002 (2002-04-04) * the whole document *	1-38	A61K39/39 C07H21/00 A61K31/70
X	WO 97 29757 A (CLEVELAND CLINIC FOUNDATION ; NAT INST HEALTH (US)) 21 August 1997 (1997-08-21) * tables 3,4 *	1-38	
X	US 5 668 265 A (CIOLKOWSKI MARY LEE ET AL) 16 September 1997 (1997-09-16) * claim 1 *	1-38	
X	US 5 495 006 A (CLIMIE SHANE ET AL) 27 February 1996 (1996-02-27) * figures 3,4 *	1-38	
P,X	WO 03 035836 A (HYBRIDON INC) 1 May 2003 (2003-05-01) * the whole document *	1-38	
P,X	WO 03 000922 A (TUCK STEPHEN F; DYNAVAX TECHNOLOGIES CORP ; DINA DINO (US); FEARON) 3 January 2003 (2003-01-03) * the whole document *	1-38	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A61K C07H
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 8 October 2003	Examiner Bardili, W
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			

EPO FORM 1503 03 02 (P04001)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 03 01 7211

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-10-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0226757	A	04-04-2002	AU 9475001 A	08-04-2002
			CA 2423487 A1	04-04-2002
			EP 1322656 A2	02-07-2003
			WO 0226757 A2	04-04-2002
			US 2002137714 A1	26-09-2002

WO 9729757	A	21-08-1997	AU 708535 B2	05-08-1999
			AU 2129897 A	02-09-1997
			CA 2246503 A1	21-08-1997
			CN 1215994 A	05-05-1999
			EP 1007655 A1	14-06-2000
			JP 2000506384 T	30-05-2000
			WO 9729757 A1	21-08-1997
			US 5998602 A	07-12-1999
			US 6214805 B1	10-04-2001

US 5668265	A	16-09-1997	US 5658738 A	19-08-1997

US 5495006	A	27-02-1996	CA 2116465 A1	02-09-1994
			AU 2657392 A	27-04-1993
			CA 2119927 A1	01-04-1993
			WO 9306122 A1	01-04-1993

WO 03035836	A	01-05-2003	WO 03035836 A2	01-05-2003
			WO 03057822 A2	17-07-2003

WO 03000922	A	03-01-2003	WO 03000922 A2	03-01-2003
			US 2003175731 A1	18-09-2003

EPO FORM P4459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)